

Research Article

Cite this article: Lincoln S *et al.* (2022). Climate change impacts on the coral reefs of the UK Overseas Territory of the Pitcairn Islands: resilience and adaptation considerations. *Journal of the Marine Biological Association of the United Kingdom* **102**, 535–549. <https://doi.org/10.1017/S002531542200090X>

Received: 14 December 2021

Revised: 30 September 2022

Accepted: 17 October 2022

Key words:

Adaptation; climate change; coral reefs; marine protected area; Pitcairn Islands; resilience

Author for correspondence:

Susana Lincoln,
E-mail: susana.lincoln@cefias.gov.uk

© Crown Copyright - Crown, 2022. Published by Cambridge University Press on behalf of Marine Biological Association of the United Kingdom. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/4.0>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.



Climate change impacts on the coral reefs of the UK Overseas Territory of the Pitcairn Islands: resilience and adaptation considerations

Susana Lincoln¹ , Benjamin Cowburn¹, Ella L. Howes^{1,2}, Silvana Birchenough¹, John K. Pinnegar¹, Stephen Dye¹, Paul Buckley¹, Charles Sheppard³, Colette C.C. Wabnitz⁴ , Leo X.C. Dutra^{5,6}, Jennifer Graham¹, Simeon Archer-Rand¹, Emily Hardman⁷, Georg H. Engelhard¹  and Bryony L. Townhill¹

¹Cefas Laboratory, International Marine Climate Change Centre, The Centre for Environment, Fisheries and Aquaculture Science, Pakefield Road, Lowestoft NR33 0HT, UK; ²Environment and Geography Department, University of York, Heslington, York YO10 5DD, UK; ³School of Life Sciences, Gibbet Hill Campus, The University of Warwick, Coventry CV4 7AL, UK; ⁴Institute for the Oceans and Fisheries, The University of British Columbia, AERL, 2202 Main Mall, Vancouver, British Columbia V6T 1Z4, Canada; ⁵CSIRO Oceans & Atmosphere, Queensland Biosciences Precinct, 306 Carmody Rd, St Lucia, QLD 4067, Australia; ⁶Centre for Marine Socioecology, University of Tasmania, Hobart, Australia and ⁷Global Marine Team, Marine Management Organisation, Hornbeam House, Electra Way, Crewe CW1 6GJ, UK

Abstract

The coral reefs of the Pitcairn Islands are in one of the most remote areas of the Pacific Ocean, and yet they are exposed to the impacts of anthropogenic climate change. The Pitcairn Islands Marine Protected Area was designated in 2016 and is one of the largest in the world, but the marine environment around these highly isolated islands remains poorly documented. Evidence collated here indicates that while the Pitcairn Islands' reefs have thus far been relatively sheltered from the effect of warming sea temperatures, there is substantial risk of future coral decalcification due to ocean acidification. The projected acceleration in the rate of sea level rise, and the reefs' exposure to risks from distant ocean swells and cold-water intrusions, add further uncertainty as to whether these islands and their reefs will continue to adapt and persist into the future. Coordinated action within the context of the Pitcairn Islands Marine Protected Area can help enhance the resilience of the reefs in the Pitcairn Islands. Options include management of other human pressures, control of invasive species and active reef interventions. More research, however, is needed in order to better assess what are the most appropriate and feasible options to protect these reefs.

Introduction

Impacts from climate change and other human activities are causing a rapid global decline of coral reefs (Wilkinson, 2008; Burke *et al.*, 2011; IPCC, 2014a, 2022). Most of the world's tropical reefs have experienced severe bleaching events in the past 30 years, with widespread bleaching occurring in 1998, 2010 and through 2014–2017 (Heron *et al.*, 2016; Sully *et al.*, 2019). Warm-water corals are projected to decline further in the coming decades as they become increasingly exposed to thermal anomalies (Van Hooidonk *et al.*, 2016; Hughes *et al.*, 2017, 2018). Ocean acidification, storminess and the compounding impact of other anthropogenic pressures further raises the risk to corals into the future (Pandolfi *et al.*, 2011; Dutra *et al.*, 2021; IPCC, 2022). Stony corals have a disproportionately important role in the marine environment; changes in species composition can dramatically impair the ecological functioning of the reef with negative, knock-on consequences to associated biological communities (Glynn, 1993; Moberg & Folke, 1999; Baker *et al.*, 2008; Fisher *et al.*, 2015). In addition to their ecological importance, coral reefs provide a range of vital services to societies and economies, including food provision, coastal protection and tourism (Moberg & Folke, 1999; IPCC, 2014a), which are particularly important in the case of small oceanic islands.

The remote Pitcairn Islands group in the equatorial South Pacific includes the islands of Pitcairn, Oeno, Henderson and Ducie, with an aggregated land area of only 49 km² (Irving & Dawson, 2012). These islands are part of an outcrop along the Foundation Seamount chain in the central South Pacific (Figure 1). Pitcairn Island is volcanic and the only inhabited island, with 52 permanent residents as of January 2020 (Pitcairn Miscellany, 2020). The coastline of Pitcairn Island is rocky and steep and exposed to large ocean swells (Avagliano *et al.*, 2016). Henderson Island is formed from a raised coral reef, and Oeno and Ducie are low coral atolls (Spencer, 1995; Robinson *et al.*, 2017). The Pitcairn Islands Marine Protected Area (MPA) was designated in 2016 (The Government of the Pitcairn Islands, 2017) and is one of the largest in the world, covering 841,910 km² (UNEP-WCMC & IUCN, 2021). The




Fig. 1. Map of the Pitcairn Islands designated Marine Protected Area, which occupies the entire Exclusive Economic Zone. The no-take MPA extends over 841,910 km², with subsistence fishing allowed within coastal conservation areas.

Pitcairn Islands MPA encompasses the entire Pitcairn Exclusive Economic Zone (Figure 1) and is a highly protected, no-take area (The Government of the Pitcairn Islands, 2017). Marine Conservation Regulations are in preparation, and in 2017 a Fisheries Management Plan was introduced, to manage the only artisanal fisheries that are allowed within coastal conservation areas, mainly around Pitcairn Island (The Government of the Pitcairn Islands, 2017; Irving *et al.*, 2019; Dawson & Irving, 2020).

The reefs of the Pitcairn Islands are located on the limits of coral growth at the eastern edge of the Indo-Pacific biogeographic realm, and are distinct from other reef systems due to their isolation. Some endemic species have been identified primarily at Ducie Island, the most remote island in the group, consisting of fish species endemic to the south-eastern tropical Pacific region as well as a few other species only found in these islands (Friedlander *et al.*, 2014). Coral reefs are present around all four of the Pitcairn Islands, with Ducie having the highest coral cover, 56%, and Pitcairn the lowest, 5% (Friedlander *et al.*, 2014). At Pitcairn Island, macroalgae are abundant (far more so than at the other three islands) and corals are predominantly absent above 10 m depth, possibly due to the combined impact of wave action and sand scouring, but are abundant below 10–30 m and deeper, due to the exceptional water clarity (Irving & Dawson, 2013; Duffy, 2014; Friedlander *et al.*, 2014).

In this study we review the risk of climate change impacts to the coral reefs of the Pitcairn Islands, together with other local human pressures, in order to support the Pitcairn Islands MPA and inform potential management and resilience actions.

Materials and methods

A literature review of climate change and coral reefs of the UK Overseas Territories of the Pitcairn Islands, as well as the British

Indian Ocean Territory (BIOT), was undertaken. This review, available as a separate technical report document (see Lincoln *et al.*, 2021), covered key drivers of climate change and observed and expected impacts on the coral reefs and other important marine life. A policy summary was also produced, to inform decision making and conservation activities in these territories (see BlueBelt Report Card, 2021). The marine environment around the Pitcairn Islands remains vastly under-studied with few published scientific studies available. The scarcity of relevant literature prevented the use of a systematic review method. Instead, the authors used the information from key published studies of the Pitcairn Islands, and mined further literature from those and from other relevant regional and global studies. Where possible, climate projections were based on either of the main Representative Concentration Pathways (RCP) climate model projections: low (mitigation) emissions scenario (RCP2.6), medium (stabilization) scenarios (RCP4.5/RCP6) and high (baseline) emission scenarios (RCP8.5; Van Vuuren *et al.*, 2011). RCP2.6 and RCP8.5 are extreme scenarios, and it is expected that actual future conditions will fall somewhere in between.

This article condenses the main findings of the original review report by Lincoln *et al.* (2021), offers a further insight into climate change risks to the coral reefs of the Pitcairn Islands, and discusses options that could help enhance the resilience of these reefs.

Results

Global Coral Reef Monitoring Network observations at Pitcairn Island between 2009–2014, found coral cover ranging from 10–20% (Moritz *et al.*, 2018). An expedition to the islands in 2012 found Ducie to have a higher live coral cover (56%) compared with Oeno (28%) and Henderson (24%) Islands, while Pitcairn Island showed the lowest percentage (5%) of live coral cover

(Friedlander *et al.*, 2014; Irving *et al.*, 2019). Reefs appeared to be mostly composed of *Porites* spp., known to be thermally resistant, as well as more sensitive *Pocillopora* and *Millepora* (Irving & Dawson, 2013; Friedlander *et al.*, 2014; Irving *et al.*, 2019).

The following sections examine in detail evidence of climate change impacts and the potential future risks to these coral reefs.

Sea temperature

SST conditions

Sea surface temperature (SST) is defined as the subsurface bulk temperature in the top few metres of the ocean measured by ships, buoys and drifters (IPCC, 2013). In the tropical Pacific Ocean, SST has increased by 0.3°C between 1950 and 2009, while in the coral reef province of the western Pacific, SST has risen by over 0.4°C in the warmest months over the same period (IPCC, 2014a). Each year, trade winds and currents push warm equatorial water eastwards, creating a 'warm pool', also known as the equatorial Pacific warm water volume (Capotondi *et al.*, 2020). This warm pool extends further east towards the Pitcairn Islands during El Niño years, and its extent has increased since the 1980s (Weller *et al.*, 2016; Sutton, 2018). As the equatorial Pacific warm pool expands, a larger area of the tropical Pacific will experience higher than average annual temperatures over the rest of the 21st century (Sutton, 2018; Capotondi *et al.*, 2020). The warm pool is controlled by the El Niño/Southern Oscillation (ENSO), but the high complexity of the ENSO phenomenon, its cyclic variability and increasing amplitude, hinder the ability of climate models to represent the diversity of El Niño events (Capotondi *et al.*, 2020). At present, it is not possible to consistently predict changes in ENSO events over the coming century (IPCC, 2013; Cai *et al.*, 2020; Capotondi *et al.*, 2020; Fang & Yu, 2020).

The Pitcairn Islands are situated along the south-eastern edge of the tropical Pacific, where heat is absorbed near the equator and then cools towards the poles (Sutton, 2018). Three intense marine heatwaves were detected in 1995, 2006 and 2017 respectively, with temperature peaks that were more noticeable at Oeno, Pitcairn and Henderson Islands (Figure 2). However, the natural variability of SST of the Pacific Ocean, which is driven by the Pacific Decadal Oscillation and ENSO system and also influenced by the Pacific warm pool, makes it difficult to attribute SST increases to climate change (IPCC, 2014a). Nevertheless, recent reports indicate that ocean conditions in the wider South Pacific region are changing faster and more acutely than initially predicted due to climate change (IPCC, 2019a; Grose *et al.*, 2020 – specifically for the region of Australia; IPCC, 2021). Bearing in mind these uncertainties, under medium emissions scenarios RCP4.5 or RCP6 (Van Vuuren *et al.*, 2011), SST around the Pitcairn Islands is projected to rise by ~0.5°C by the middle of this century and by 0.9°C by the end of this century (NOAA Climate Change Portal, 2020). Under a high emissions scenario RCP8.5 (Van Vuuren *et al.*, 2011) and by the end of this century, average SST is expected to warm by 2–2.5°C (Sellar *et al.*, 2019).

Degree Heating Weeks (DHW) or Degree Heating Months (DHM) represent the number of weeks or months the corals are exposed to warming above a given threshold multiplied by temperature above the threshold. Based on SST satellite data, peaks of thermal stress expressed as DHWs occurred in 1995, 2006 and 2017 (see Figure 2 and Table 1), with Oeno reaching Bleaching Alert 1 (DHW 4–8) in all those three years (NOAA Coral Reef Watch, 2018). Pitcairn Island was the only island to experience DHWs higher than 8 (Bleaching Alert 2) during 2006, while Ducie Island did not experience thermal stress enough to trigger a bleaching alert (DHW less than 4; see Figure 2 and Table 1; NOAA Coral Reef Watch, 2018).

While there is no clear evidence of coral bleaching from sea-water warming in the Pitcairn Islands, there are observations of coral damage caused by unusual oceanic cold intrusions in the Ducie atoll, with up to 40% coral damage by cold-water stress found between 10–15 m depth, as explained further in the extreme weather events section (Irving & Dawson, 2013; Dawson & Zhang, 2020). However, without baseline data it is not possible to accurately assess these factors or changes over time. Significant acute thermal stress (DHW 8.9) was experienced at Pitcairn Island in 2006, and Oeno Island (DHW 6) in 2006 and 2017, but otherwise the risk of acute thermal stress in the archipelago appears to remain low (Table 1). Undetected episodes of coral bleaching and mortality over the past 35 years in the Pitcairn Islands, however, remain a possibility.

Future SST projections

The Pitcairn Islands are located near the southern limit for optimal reef growth, with Ducie being the most southerly atoll island in the world. As a result of cooler conditions and slow projected warming, future SST conditions are likely to remain relatively suitable for coral growth, at least until the middle of this century (Couce *et al.*, 2013; Jones *et al.*, 2019). In the Pitcairn Islands and under a high emissions scenario, the onset of severe annual bleaching events is not expected until 2059, which is later than the average global onset threshold of 2043 (Van Hooidonk *et al.*, 2016; Sellar *et al.*, 2019). In addition, most reefs in the Pitcairn Islands are below 10 m depth (Irving & Dawson, 2013; Duffy, 2014; Friedlander *et al.*, 2014), with the exception of the shallow reef flats at Oeno and Henderson and those in the Ducie lagoon, and it is expected that those deeper fore-reefs will be relatively sheltered from thermal stress (Nakamura & Van Woosik, 2001; Muir *et al.*, 2017; Cowburn *et al.*, 2019). Deeper areas of the outer-edge reefs on the Great Barrier Reef have been found to provide refuge from bleaching for *Acropora*, *Pocillopora* and *Porites* spp. (Baird *et al.*, 2018). In the eastern Pacific, a depth refugia has also been demonstrated for *Millepora* spp. (Smith *et al.*, 2014). However, the efficacy of these deeper reefs as refugia from warming will be relative, and will depend on the severity of the warming events and on the thermal sensitivity of the corals, as other evidence from the wider Pacific and from the Caribbean shows (Smith *et al.*, 2016; Venegas *et al.*, 2019). The extreme clarity of the water in the Pitcairn Islands also allows for deeper irradiance, which could to some extent counteract the cooling effect at depth. At Pitcairn Island, corals grow on reefs that reach down to 75 m depth and are dominated by thermally resistant *Porites* spp. and *Pocillopora* sp. and the hydrocoral, *Millepora* spp., which are more likely to withstand warming conditions (Irving & Dawson, 2013; Friedlander *et al.*, 2014). However, in similar communities in the Central and East Pacific, those same coral species were found to die in large numbers during bleaching events (Glynn, 1984; Pratchett *et al.*, 2013). This suggests that the corals of the Pitcairn Islands would still be relatively susceptible to thermal stress (Figure 2; Table 1), particularly in the shallower reefs around Ducie Island dominated by more sensitive corals such as *Acropora* and *Montipora* spp. (Loya *et al.*, 2001).

Ocean acidification

Surface pH and aragonite saturation conditions

The increasing uptake of CO₂ by the world's oceans is manifested as changes to ocean carbon chemistry particularly pH and saturation state of calcium carbonate minerals such as aragonite and calcite, conditions which are critical to the growth and survival of corals and other calcifying organisms.




Fig. 2 Plots of thermal stress in the Pitcairn Islands showing: Time series of annual total Degree Heating Weeks (DHWs) from the four main islands (top right); time series of mean annual sea surface temperature (SST) (top left); and maps of DHWs during strong thermal stress years 1995, 2006, and 2017, showing the position of the islands. The dark blue areas denote no data. Data obtained from NOAA Coral Reef Watch (2018).

The largest changes in pH are projected for the surface layer except in the subtropics, where they are expected to occur where carbonate buffering capacity is lower between 200 and 300 m (IPCC, 2013). There is no long-term surface water pH data for the equatorial Pacific, making it difficult to ascertain trends (Lenton et al., 2018; GOA-ON, 2020). A range of surface pH 7.91–8.12 has been previously estimated for the equatorial Pacific between 1997–2011 (Sutton et al., 2014), values that are lower, and appear to be declining faster, than other open-ocean regions (Johnson et al., 2015; Lauvset et al., 2015; Wagener et al., 2018). The pH of the wider tropical Pacific is projected to decrease by 0.15 pH units by 2050 under a high emissions scenario (Johnson et al., 2015). Around the Pitcairn Islands, mean surface water pH is currently estimated near 8.12 and by the

end of this century could decrease between 0.08 and 0.27 units, according to low and high emissions scenarios respectively (Figure 3; NOAA Climate Change Portal, 2020).

The saturation states of calcium carbonate minerals naturally decrease with depth as total dissolved CO₂ increases from a combination of respiration and cold temperatures, and there are indications that surface ocean aragonite saturation state in the Pacific has already declined from 4.5 to 3.8 (Feely et al., 2009). Aragonite saturation below 3 is considered extremely marginal for coral growth, and no major reefs currently exist at such conditions (Langdon & Atkinson, 2005; Lenton et al., 2018); while unprotected shells and skeletons will begin to dissolve when surrounding seawater saturation level declines below 1 (Feely et al., 2009; Birchenough et al., 2017). Levels of aragonite saturation in the South Pacific are already thought to be close to the point where it can weaken coral skeletons and slow down coral growth (Dutra et al., 2018). However, direct field observations linking acidification and ocean chemistry changes to physiological impacts on corals are still rare (Doney et al., 2020; Doo et al., 2020). It is possible that the responses of corals to more acidic waters are still within the ranges of natural variability, and such responses would also be influenced by local conditions of water quality and circulation patterns (Field et al., 2014). There are hardly any *in situ* studies of ocean acidification effects on corals in Pacific islands (Dutra et al., 2018), while studies of ocean acidification impacts in other regions have found alterations in the physiology of the corals and the wider ecology of the reef, from changes in the pH of the calcifying fluid of the corals (Kubota et al., 2017), to loss of skeletal density (Mollica et al., 2018), to shifts in the species composition of colonies as well as increased algal cover and bioerosion (Fabricius et al., 2011; Enochs et al., 2015, 2016), to loss of ecological complexity and function of the reef (Mumby, 2009; Anthony et al., 2011).

Table 1. Past thermal stress in each of the Pitcairn Islands showing mean SST at the start and end of a time series spanning 1985–2018 and the difference in mean SST, along with Degree Heating Weeks (DHWs) in potential bleaching years, the cumulative DHW, and the number of Alert Level 1 and 2 thermal stress events. Data obtained from NOAA Coral Reef Watch (2018).

Parameters of thermal stress	Oeno	Pitcairn	Henderson	Ducie
Mean SST (1985–89)	24.79°C	24.32°C	24.5°C	24.16°C
Mean SST (2013–18)	24.82°C	24.19°C	24.55°C	24.31°C
Difference in mean SST	0.03°C	−0.13°C	0.05°C	0.15°C
DHW 1995	4.6	3.1	1	1.1
DHW 2006	5.8	8.9	5.5	2.5
DHW 2017	6	3.6	2.8	0.5
Cumulative DHW	16.4	15.6	9.3	4.1
Alert 1 (DHW 4–8) frequency	3	2	1	0
Alert 2 (DHW >8) frequency	0	1	0	0

Future pH and aragonite saturation projections

ENSO dominates ocean climate in the Pacific, including ocean acidification and ocean carbon chemistry: during non-El Niño




Fig. 3. Outputs from the CMIP5 ensemble showing mean surface water pH in the eastern Indian Ocean for the period 1956–2005. Image provided by the NOAA/OAR/ESRL Physical Sciences Laboratory, Boulder, CO, USA, from their website at <http://psl.noaa.gov/> (NOAA Climate Change Web Portal, 2020). The approximate location of the Pitcairn Islands is indicated by the orange star symbol.

periods, ocean acidification has been observed to progress more rapidly due to the influx of upwelling deep, CO₂-rich water (McPhaden *et al.*, 2006; Sutton *et al.*, 2014; Fiedler & Lavín, 2017). Projections of aragonite saturation state to the end of this century under high emissions scenarios for the central South Pacific surrounding the Pitcairn Islands suggest that it is likely to decrease to 2.5 (Sellar *et al.*, 2019, see Figure 4), a sub-optimal level for coral calcification. But the confidence in these projections is only limited, due to the variability of the ENSO and the associated upwelling forcing, and the influence that these processes have on future ocean pH levels in the Pacific Ocean (Christensen *et al.*, 2013 and references therein). In

addition, saturation of calcium carbonate minerals and pH can also vary at local scales depending on the habitat mosaic present (Cyronak *et al.*, 2018; Pacella *et al.*, 2018; Bergstrom *et al.*, 2019; Liu *et al.*, 2020) and it can be affected by biological processes like photosynthesis and respiration (Hurd *et al.*, 2018; Fabricius *et al.*, 2020).

The South Pacific Circulation Gyre is expected to slow down future changes in carbonate saturation and pH in the Pitcairn Islands region compared with the wider oceanic basin (Hoegh-Guldberg & Bruno, 2010; Friedlander *et al.*, 2014). However, across the tropical Pacific by mid-century under a high emissions scenario, average aragonite saturation is likely to




Fig. 4. Average sea surface aragonite saturation state (Ω_{Arag}) for the Pitcairn Islands, showing the predicted decline in saturation under RCP2.6 (left) and RCP8.5 (right). Data from the UKESM1 forecast model, from Sellar *et al.* (2019).

decline to near 3, meaning a shift to sub-optimal conditions (Johnson *et al.*, 2015). This will cause a decline in coral calcification rate of about 10% (Chan & Connolly, 2013), which is projected to worsen as saturation continues to decline towards the end of the century.

Sea level rise and extreme weather events

Observed sea level rise and storminess trends

Sea level has risen across the Pacific Ocean by varying amounts due to large-scale climate processes, and extreme sea levels have also been detected driven by a combination of long-term sea level rise from climate change and short-term factors such as extreme tides, storm surges and the associated wind-wave setup (Australian Bureau of Meteorology and CSIRO, 2014). There are regional differences in sea level rise across the tropical Pacific (PSMSL, 2020), with the highest rates, ~ 12 mm year⁻¹, detected in the western South Pacific between 1993–2009 (Meyssignac *et al.*, 2012; Nurse *et al.*, 2014). These high sea level rise rates are thought to reflect modulations by natural phenomena such as ENSO, with lower/higher-than-average variability of sea levels during El Niño/La Niña events of the order of ± 20 to 30 cm in the western tropical Pacific (Cazenave & Remy, 2011; Becker *et al.*, 2012). There are no tide gauge records from the Pitcairn Islands (Woodworth & Hibbert, 2015) but indirect estimations suggest that sea level has already risen almost 4 times the global average. It is thought that the low-lying atolls of Ducie and Oeno have persisted, and even increased in surface area and elevation by way of seasonal erosion and accretion processes following extreme events and changes in sediment supply (McLean & Kench, 2015). This has also been confirmed for other small atoll islands in the South Pacific, where high sea level rise rates have, as yet, not caused loss of dry land area overall (Kench *et al.*, 2015).

Storms can inflict substantial damage to shallow coral reefs. In the Pacific Ocean, the intensity of tropical cyclones appears to have increased since 1990 (Diamond *et al.*, 2013), although it is not clear whether there is a basin-wide trend in the number of very intense tropical cyclones across the Pacific basin (Australian Bureau of Meteorology and CSIRO, 2014; Holland & Bruyère, 2014; CSIRO *et al.*, 2015; Hoarau *et al.*, 2018; Lafale *et al.*, 2018; Duthiel *et al.*, 2020), and the attribution of this trend to climate change also remains in question (Collins *et al.*, 2019; Chand *et al.*, 2020). The trajectory and frequency of tropical cyclones in the South Pacific is typically driven by the natural variation of ENSO (Lafale *et al.*, 2018), and severe storms are more likely linked to extreme El Niño events with an extensive eastward propagation (Santoso *et al.*, 2013; Stephens & Ramsay, 2014). Deep ocean swells generated by extratropical cyclones in mid and high latitudes are also known to occasionally reach Pacific islands (IPCC, 2014b). For example, oceanic swells originated in the North Pacific, especially during years with a strong El Niño, have caused damage to reefs in Hawaii (Rooney *et al.*, 2004; Fletcher *et al.*, 2008). At Henderson Island, sections of fore-reef have been eroded exposing the bedrock underneath and coral rubble has been found below 30 m depth, all of which is likely to have been caused by storm wave action (Irving & Dawson, 2013). At Ducie Island, there are observations of coral damage caused by unusual oceanic cold intrusions in the atoll. Rehder & Randall (1975) documented up to 40% coral damage by cold-water stress found between 10–15 m depth recorded during an expedition in 1970. During diver surveys between 2016–2018, coral bleaching was documented between 10–15 m depth and linked to cold-water anomalies following the 2015–2016 El Niño event, confirmed by SST satellite data (Rehder & Randall, 1975; Irving & Dawson, 2013; Dawson & Zhang, 2020). Also in Ducie Island

in 1975, parts of the land were found deforested by high waves (Rehder & Randall, 1975). There is also evidence of transient conditions of coastal turbidity and sediment runoff at Pitcairn Island in 2012 following a major rainfall event, the heaviest since records began, although it is thought that the prevailing strong currents around the island would have dispersed the suspended sediments, mitigating the effects of turbidity and siltation (Irving & Dawson, 2013; Friedlander *et al.*, 2014; Irving *et al.*, 2019). Given the small size of Pitcairn Island and its population and the domestic scale of current land use practices, the low cover of live coral could be attributed to exposure to wave action, rather than as an effect of runoff and sedimentation.

Future sea level rise and storm projections

Sea levels are highly variable in the Pacific, but it is very likely that under high emissions scenarios and by the end of this century, the South Pacific in particular will experience the largest sea level rise globally, ranging 54–217 cm (Nurse *et al.*, 2014; Lafale *et al.*, 2018; Vousdoukas *et al.*, 2018). This presents a significant risk for small coral islands, especially when combined with a potential intensification of tropical cyclones and degraded fringe reefs (The World Bank, 2016; Lafale *et al.*, 2018). A rise of more than 1 m by the end of the century is considered to be equivalent to a one in 100-year extreme sea level event currently (Vousdoukas *et al.*, 2018; Kench *et al.*, 2018a; Masselink *et al.*, 2020; Tajima *et al.*, 2020). For the Pitcairn Islands however, sea level projections are still highly uncertain (Zhang *et al.*, 2014) due to lack of resolution and long-term data, but there is an expectation that the rate of sea level rise is likely to accelerate and threaten the smaller low-lying islands (Kench *et al.*, 2015), particularly as compounded with coral decalcification.

Mean significant wave heights appear to have increased in regions of the Pacific and Southern Oceans, but there is as yet low confidence in these projections and the lack of data means it is challenging to discern between long-term climate trends and natural variability (Wang & Swail, 2006; Young *et al.*, 2011; Seneviratne *et al.*, 2012; Church *et al.*, 2013). At present however, most tropical cyclones in the South Pacific basin are formed in the region of the Vanuatu archipelago (Diamond *et al.*, 2013), strengthening as they move southwards, and so the Pitcairn Islands tend to miss the most severe events (Lafale *et al.*, 2018). Instead, the impact of long-distance ocean swells from remote low-pressure systems such as the Southern Ocean could be more significant for the Pitcairn Island corals.

Other human pressures in the Pitcairn Islands

In spite of their isolation and the small size of the resident human community, non-climatic anthropogenic pressures – some trans-boundary – may be impacting the marine environment of the Pitcairn Islands, and particularly the corals.

Fishing

Prior to 2016, most non-commercial fishing took place around Pitcairn Island, with occasional (every 4 to 5 years) trips to 40-Mile Reef to the east-south-east of the island (Irving and Dawson, 2012). There is no mandatory recording of catches at present and only intermittent fisheries monitoring, but reconstructed data indicate that landings by this small-scale fishery have been higher than the data reported by the Food and Agriculture Organization of the United Nations on behalf of the Pitcairn Islands, although they declined to 4 tonnes year⁻¹ overall by 2014 (Sea Around Us, 2016; Coghlan *et al.*, 2017). Licenced commercial fishing declined sharply after 2005 as fleets withdrew to international waters outside the EEZ (Irving & Dawson, 2012). All forms of fishing have been banned within

the Pitcairn Islands' MPA since its designation in 2016, apart from artisanal fishing by the local community which is allowed within the coastal conservation areas. There have been some concerns of overfishing targeting groupers, sharks and lobsters (Irving & Dawson, 2012; Coghlan *et al.*, 2017) which could have negative consequences for the reef since some of these species, sharks particularly, play an important role in its resilience (Roff *et al.*, 2016). A Fisheries Management Plan was introduced in 2017 for the management of the local fishery, particularly around Pitcairn Island (Irving *et al.*, 2019; Dawson & Irving, 2020). The management plan brought in minimum catch sizes for a number of species of fish and shellfish and also banned the fishing of sharks.

Marine litter

Ocean-borne litter is an example of transboundary pressure in the Pitcairn Islands. Henderson Island is reported to have one of the highest levels of marine plastic litter on the planet despite it being uninhabited (Lavers & Bond, 2017; Forrest & Hindell, 2018; Irving *et al.*, 2019; Ryan, 2020; Ryan & Schofield, 2020). This island is on the edge of the South Pacific Gyre, which collects litter from distant Pacific shores and deposits it on the island's shores (UK Gov Marine Developments, 2019). On the beaches of Henderson below the accumulated plastic, temperature within the sand was found to be +2.5°C above the daily maximum and dropped down by -1.5°C below the daily minimum, compared with areas with no plastic debris (Lavers *et al.*, 2021). Quantities of plastic items were also found tangled on the surrounding reefs (S. Archer-Rand, personal communication).

Anchoring

Anchoring from visiting cruise ships and yachts and from supply vessels is another concern (Irving *et al.*, 2019). Anchors and anchor warps can destroy large swathes of seabed as they settle and when retrieved (Giglio *et al.*, 2017). Hard corals are especially susceptible to the impacts of anchoring, but the scale of damage and recovery timeframes remains unknown.

Non-native species

The presence of invasive non-native species, even on land, as well as coral diseases, remain important threats to coral reefs. Invasive rats on Henderson Island have caused dramatic declines of the biomass of resident seabirds, including endemic or endangered species (BirdLife International, 2020) and this could cause an indirect negative effect on the surrounding reefs as found in other oceanic coral islands, where declines in nesting seabird populations caused losses in reef productivity due to the loss of guano nutrient inputs (Graham *et al.*, 2017, 2018; Benkwitt *et al.*, 2019, 2021). The eradication of invasive rats should be important for the sustainable management of the natural environment of these islands (Graham *et al.*, 2017; Benkwitt *et al.*, 2019; BirdLife International, 2020; Perez-Correa *et al.*, 2020). Corallivorous crown-of-thorns starfish have been reported in the Pitcairn Islands (Birkeland, 1989; Irving, 1995), as well as coral diseases.

Discussion

Corals can recover following bleaching and other climate change impacts such as storm damage, and healthy and well-balanced reefs hold the key to this resilience. Herbivorous reef-fish will for example encourage recruitment of new corals by grazing on the algae and preventing their encroachment on the reef (Mumby *et al.*, 2016; Steneck *et al.*, 2019). Good water quality is also particularly important to help recovery following bleaching and to reduce susceptibility to disease outbreaks and invasive

species (Carlson *et al.*, 2019; MacNeil *et al.*, 2019). Large MPAs are effective options for reef resilience, as they minimize local stresses and support a wide range of species and functional groups, particularly when they include replicates of representative habitats, ensuring reef connectivity, and encompass different thermal regimes (Carpenter *et al.*, 2008; McLeod *et al.*, 2019). In the Pitcairn Islands MPA in particular, the deeper, cooler fore-reefs could act as areas of refugia from the effects of warming to corals and other species (Kleypas *et al.*, 2010; Friedlander *et al.*, 2014).

This study suggests that, in the Pitcairn Islands, thermal stress is likely to remain low compared with other tropical reefs globally, particularly in the deeper fore-reefs, although those deeper coral communities appear dominated by sensitive species which may heighten their vulnerability to thermal stress should it occur. The geographic isolation of these reefs is another challenge in terms of supply of healthy larvae, meaning these reefs would rely on self-seeding to boost recolonization and recovery (Robinson *et al.*, 2017; Romero-Torres *et al.*, 2018). On the other hand, the low level of other local detrimental pressures in these islands such as nutrient inputs or pollution, can help corals better withstand thermal stress (Wiedenmann *et al.*, 2013). The additional protection granted by the MPA is expected to promote recovery following bleaching (Sheppard *et al.*, 2008; Gilmour *et al.*, 2013), although there is evidence that severe bleaching and mortality can still occur in highly protected reefs, as demonstrated by the Aldabra Atoll in the Western Indian Ocean (Cerutti *et al.*, 2020), BIOT (Head *et al.*, 2019), Jarvis Island in the central Equatorial Pacific (Vargas-Ángel *et al.*, 2019) and the Great Barrier Reef (Hughes *et al.*, 2017).

There are two main approaches for enhancing climate resilience of ecosystems: management of local pressures and implementation of adaptation and resilience-building actions. Efforts are increasing globally to understand coral recovery and promote adaptation using coordinated ecosystem-based management approaches (Obura & Grimsditch, 2009; NOAA Coral Reef Conservation Programme, 2014; Belokurov *et al.*, 2016; Harvey *et al.*, 2018; Roche *et al.*, 2018; McLeod *et al.*, 2019). The goal of these adaptation strategies is to 'maintain the ecosystem in a healthy, productive, and resilient condition so that it can continue to provide the services that society wants and needs' (NOAA Coral Reef Conservation Programme, 2014).

Here, we discuss potential options for enhancing the climate resilience of coral reefs in the Pitcairn Islands, based on examples from other reefs. Intervention and management actions should be underpinned by scientific evidence wherever possible, and include monitoring plans, review periods and adaptive management approaches (Belokurov *et al.*, 2016). Decisions and actions should also be undertaken in consultation with and with the full involvement of the local community, as custodians and direct beneficiaries of these reefs, so they can contribute their valuable knowledge. An example of a step-by-step process that could be followed to develop an Action Plan for building and enhancing climate change adaptation and resilience of the coral reefs of the Pitcairn Islands is represented in Figure 5.

Next, we discuss a number of considerations to develop a plan of climate adaptation and resilience for the Pitcairn Islands' reefs, that could potentially be incorporated into the framework of the Pitcairn Islands MPA.

Enhancing resilience through management of local pressures

Reducing pressures from human activities and regulating the uses of the marine environment may not entirely prevent coral bleaching, or coral mortality, but boost the resilience of reefs and their chances of recovery (Nyström *et al.*, 2008; IPCC, 2014a; UNEP, 2017). Approaches of this kind can help delay the onset of



Fig. 5. Conceptual diagram for the development of a climate change adaptation and resilience action plan for the coral reefs of the Pitcairn Islands, adapted from Obura & Grimsditch (2009) and Belokurov *et al.* (2016).

mass loss of corals, allowing time for emissions mitigation measures to take effect in limiting global warming (IPCC, 2014a, 2019b). There are strict regulations as part of the designation of the Pitcairn Islands MPA, a highly protected marine area and one of the largest in the world at 841,910 km² (UNEP-WCMC & IUCN, 2021). They include a 5-year Management Plan to limit local pressures as well as long-term monitoring of the coral reefs to assess reef health. This vast MPA hosts a network of shallow and deep reefs across a range of different thermal regimes, likely to offer taxonomic overlap and a well-connected reef system with areas of refugia and therefore appears to meet reef resilience management requirements (McLeod *et al.*, 2019). Given their isolated location and low population, the Pitcairn Islands are exposed to fewer pressures compared with other similar coral atoll islands, but nevertheless the following pressures have been identified in the Pitcairn MPA Management Plan: Illegal, Unreported and Unregulated (IUU) fishing by foreign fishing vessels; pollution including oil spills and marine plastics; anchor damage; erosion and run-off from land; and invasive non-native species.

Fishing

The implementation of large, highly protected marine areas such as the Pitcairn Islands MPA can be an effective climate resilience option for the corals and other marine communities (Hughes *et al.*, 2003; McClanahan *et al.*, 2008; Coghlan *et al.*, 2017; Hays *et al.*, 2020). The Pitcairn Islands MPA limits local fishing activities to coastal conservation areas (Pitcairn Government, 2017). A Fisheries Management Plan regulates the inshore artisanal fishery including gear restrictions and minimum landing sizes (Irving *et al.*, 2019; Dawson & Irving, 2020), and a separate Compliance and Enforcement Strategy covers the surveillance and prosecution of IUU fishing offences. An immediate ecological benefit to the reef results from allowing healthy stocks of herbivorous reef-fish and invertebrates to remain and graze on the algae that otherwise could occupy the reef, freeing up substrate for coral recruitment (Mumby *et al.*, 2007; Steneck *et al.*, 2019). There is evidence in the Caribbean region, where measures such as this enabled recovery of damaged reefs following hurricane impacts and bleaching (Steneck *et al.*, 2019).

Predatory fish are also important to keep smaller, corallivorous fish, and other trophic groups in check and therefore preserve a healthy functional reef. Previous fish assessments indicate that whilst fish communities in the Pitcairn Islands are relatively healthy, numbers of sharks and groupers have been locally impoverished due to targeted catches (Friedlander *et al.*, 2014; Coghlan *et al.*, 2017; Dawson & Irving, 2020; Duffy *et al.*, 2021).

Pollution and litter

Improving water quality can also promote climate resilience in corals. Ridge-to-reef coastal management approaches help control and reduce land sources of pollution and disturbance (Carlson

et al., 2019). Moreover, coral monitoring in the Great Barrier Reef show that improving water quality helps coral recovery following bleaching and it also reduces susceptibility to crown-of-thorns starfish outbreaks and diseases (MacNeil *et al.*, 2019). On Pitcairn Island, macroalgae are more abundant here compared with the other three islands, possibly as a combination of runoff (Irving *et al.*, 2017) and wave action. Changes in land use and soil erosion issues on Pitcairn Island have led to excess runoff inputs into coastal waters during heavy rain events resulting in short-term sedimentation turbidity (Friedlander *et al.*, 2014). However, coastal waters around all four islands are unusually clear overall suggesting oligotrophic conditions with low concentrations of dissolved inorganic nutrients. Preserving land vegetation and encouraging sustainable agricultural practices would help prevent soil erosion and maintain good water quality.

A more noticeable issue in terms of marine pollution in the case of the Pitcairn Islands is plastic litter, with Henderson Island declared to have some of the highest levels of ocean-borne plastic litter on the planet (Forrest & Hindell, 2018; Irving *et al.*, 2019; Ryan, 2020; Ryan & Schofield, 2020). Marine plastics are known to endanger many marine animals and although a specific direct link to the corals themselves (other than by physical damage from large-size items such as entangled fishing nets) is not overly clear, it has been suggested that the proliferation of plastic litter increases the risk of coral diseases (Lamb *et al.*, 2018).

Invasive species and coral diseases

Other important threats to corals are diseases and the presence of invasive and non-native species, even on land. A dramatic example is the introduction of rats to small islands, not just in the Pitcairn Islands but also in the British Indian Ocean Territory. As the rats caused a crash in the population of breeding seabirds, this resulted in a detrimental knock-on effect to coastal reefs following the loss of nutrients inputs from the birds' guano (Graham *et al.*, 2017, 2018; Benkwitt *et al.*, 2019). Eradication of rats should therefore be a conservation priority for the islands as it could have positive effects on the nearshore corals if seabird populations are able to re-establish their numbers to previous levels (BirdLife International, 2020), and the inputs of guano nutrients to the reef areas are restored (Graham *et al.*, 2017; Benkwitt *et al.*, 2019; BirdLife International, 2020; Perez-Correa *et al.*, 2020). Finally, although very challenging given the vast size and remoteness of the Pitcairn Islands MPA, early warning, monitoring or screening surveys (for example, by applying novel techniques such as detection of environmental DNA) could help detect outbreaks of crown-of-thorns and coral diseases, both of which can have equally devastating consequences for reefs (Birkeland, 1989; Irving, 1995; Galloway *et al.*, 2009).

Anchor damage

Anchoring can damage corals and other benthic communities (Irving *et al.*, 2019). Some measures have been taken towards

limiting anchoring of cruise ships, yachts and supply vessels to certain areas, but the seabed around these locations should be monitored regularly, so that further measures can be put in place if necessary (Lewis *et al.*, 2017; Sheppard & Sheppard, 2019). For example, permanent moorings can help reduce the impacts of anchoring.

Promoting coral recovery through adaptation and restoration

Reducing future global greenhouse gas emissions is, at present, the only feasible pathway to controlling future global warming and therefore effectively limiting climate change impacts on coral reefs (Bruno *et al.*, 2019). Tangible and targeted management measures and regular monitoring can help in determining what, if any, active steps are required to further promote coral resilience and adaptation. These measures are likely to involve an array of interventions, from improving protection through MPA management, to reducing local pressures, and to restoring degraded and damaged reefs (Abelson, 2020).

Recent coral planting and seeding initiatives have had some local success in remediating coral reefs elsewhere (Doropoulos *et al.*, 2019; IPCC, 2019b). Planting or seeding of corals can draw on thermally resistant species or strains (Dance, 2019), and so while this might change the diversity or species composition of the corals in an area, it can enhance the overall resilience of the reef. Coral recruitment can also be enhanced by collecting gametes from the sea during spawning, and following fertilization returning settled larvae to the sea a few days later (Dance, 2019). Research is underway in Australia to breed heat-tolerant coral species and strains (Cornwall, 2019). Coral husbandry also allows testing if corals can adapt to warming temperatures, and genetic manipulation of the bacteria or zooxanthellae in the corals, or even the corals themselves, may allow the creation of new heat-tolerant mixtures (Crabbe, 2019). However, coral restoration is usually only targeted at reefs that reach a critical stage of damage, to aid recovery, which is not the case in the Pitcairn Islands at this moment. Furthermore, coral restoration interventions are expensive and labour-intensive, and would be logistically challenging on remote reefs as these. Reducing bleaching during heatwaves by using artificial shading at times of high sea temperatures has also been proposed (Belokurov *et al.*, 2016); again, this may not be a practical choice of intervention in the Pitcairn Islands' reefs.

All these potential interventions can be very labour intensive and costly, and often involve complex decisions (Abelson, 2020; Anthony *et al.*, 2020) and are not without uncertainties and controversy as to their efficacy. Their implementation across large and remote areas such as the Pitcairn Islands MPA presents big challenges.

Further options for coral action on the Pitcairn Islands

Monitoring the reefs and other vulnerable benthic habitats around Pitcairn, Henderson, Oeno and Ducie Islands would help identify any impacts. It would be important to adopt a strategic approach to any monitoring, optimized for example by gathering additional data to support the Fisheries Management Plan, and by coordinating efforts and maximizing coverage of the MPA, and this is already contemplated as part of a Monitoring and Research Plan associated to the Pitcairn Islands MPA Management Plan.

Knowledge gaps and future research needs

Habitat mapping and reef vulnerability assessments

There remain large data gaps on the coral communities of the Pitcairn Islands (Friedlander *et al.*, 2014) making it difficult to determine a pre-bleaching baseline, and therefore habitat

mapping of coral reefs and other benthic communities across the MPA appears to be an important and necessary step. Regular coral monitoring of key ecological indicators could also be implemented, including, but not limited to: coral cover; species composition; disease screenings; occurrence of non-native invasive species such as crown-of-thorns; coral spawning and recruitment studies; water quality; changes to macroalgal cover; and fish groups counts. A smart and targeted approach to data acquisition would ensure that the information is ecologically relevant and provide the necessary evidence to support adaptive management measures.

A coral vulnerability assessment (Belokurov *et al.*, 2016; UNEP, 2017) would be useful to characterize exposure and sensitivity to climate change or other pressures across the MPA and inform decisions on interventions. Coral monitoring can also be used to identify more resistant species or colonies (Belokurov *et al.*, 2016), so they can be prioritized for protection and research.

Finally, further investigations on the deeper reefs looking at factors such as temperature profiles, light attenuation, coral community composition, and connectivity to shallow reefs, would help determine their potential as climate refugia.

Coral conservation interventions

While some of the emerging intervention techniques such as planting, seeding and genetic engineering of coral and zooxanthellae strains may appear promising in other global reef regions, some of those methods are still debatable and require further research and at this point would be too labour-intensive and expensive to be practical or even necessary in the case of the Pitcairn Islands. It has been proposed that some of the healthier reefs in the Pitcairn Islands have the potential to become coral seed banks for use in coral conservation and restoration projects in other degraded reefs in the wider Pacific. Given the remoteness of the Pitcairn Islands, this would present important logistical challenges; however as trials in the Great Barrier Reef show it remains a possibility. In any case it would require further research, not the least because unlike corals from the Red Sea and the Persian or Arabian Gulf, which are exposed to higher levels of environmental stress, the Pitcairn Islands corals are likely to be thermally naïve and therefore sensitive.

Ocean acidification

It is important to improve our understanding of ocean carbon chemistry and coral adaptation in the Pitcairn Islands, given the high and immediate risk posed by ocean acidification. These efforts could be integrated as part of the Global Ocean Acidification Observing Network (GOA-ON) monitoring network expansion, ideally by encouraging capacity in location. Given that this is a data-poor region, and taking into consideration the potential severity of calcium carbonate minerals saturation suggested by future projections, ocean acidification research in the Pitcairn Islands deserves urgent attention. The use of autonomous sensor technology could be a feasible, cost-effective option worth considering.

Research of ocean acidification on corals focuses on single-species responses. Increasingly, experiments now combine multiple climatic and non-climatic stressors on single species, which are revealing how other stressors interact with ocean acidification. It is not yet clear to what extent corals and other marine organisms can acclimatize and adapt to ocean acidification, however. *In situ* studies are required to build an understanding of effects in single species, and extrapolate these responses to impacts on whole reef systems and functions, including food webs and competition.

Sea level

It is important to better understand the processes by which coral islands adjust to sea level rise (Barnett & O'Neill, 2012; Kench *et al.*, 2015). Rising sea level is expected to cause coastal erosion, remobilize sediments and destabilize islands, but there is debate about the likely physical changes to atolls in the future (Kench *et al.*, 2015, 2018b; Storlazzi *et al.*, 2018; Ford *et al.*, 2020). Further research into atoll coastal processes in response to sea level rise and reef erosion in the Pitcairn Islands, particularly at Ducie, Henderson and Oeno, would be important.

Implementation of fisheries management and collection of catch records

There is a need for an effective management of the local artisanal fishery (Bell *et al.*, 2011; Coghlan *et al.*, 2017; Dawson & Irving, 2020). Data and information gathered as part of the fisheries management and catch recording should be integrated and coordinated with marine monitoring activities across the wider MPA, including corals. There are plans to address this as part of the MPA Management Plan and associated Monitoring and Research Plan.

Summary and conclusions

The Pitcairn Islands are some of the most remote and best preserved oceanic island environments (Readman *et al.*, 2013; Moritz *et al.*, 2018) where the low level of other human pressures could allow them to persist longer into the future (UNEP, 2017). To date, no significant changes in average SST have been detected, although marine heatwaves have been observed (in 1995, 2006 and 2017). There is, so far, no evidence of coral bleaching from seawater warming in the Pitcairn Islands, but the reefs' species composition suggests that they would likely be sensitive to thermal stress should that occur. Also, the geographic isolation of these reefs is an added challenge in terms of larval recruitment. Although recruitment of juvenile corals has yet to be assessed, the isolation of these islands, being further than 100 km away from each other and even further removed from any other reefs, indicates limited connectivity and high reliance on self-seeding (Swearer *et al.*, 1999; Wood *et al.*, 2014). Furthermore, even if these reefs persist their composition and structural complexity is likely to change (Sheppard *et al.*, 2020). Ocean acidification appears to be a significant and more immediate risk to these reefs, likely to lead to coral de-calcification by the end of the century under a high emissions scenario.

Future climate change is also likely to bring about a host of combined pressures, feedbacks and indirect impacts on coral reefs, some of which are unknown. The knowledge and appreciation of such cumulative impacts are growing rapidly (Ateweberhan *et al.*, 2013; Cinner *et al.*, 2016; Bruno *et al.*, 2019). Even among the direct impacts, there are major differences in the understanding and confidence surrounding impact pathways, that are partly attributable to differences in the timing of impacts: for example while impacts from warming SST and storms are already being observed, those from sea level rise and ocean acidification will take time to emerge (Cinner *et al.*, 2016). As research in these fields continues, it is expected that new impact pathways will be found (Cinner *et al.*, 2016). For example, ocean acidification is likely to increase susceptibility of corals to storm damage because a lower seawater pH will weaken coral skeleton density and strength (Hoegh-Guldberg *et al.*, 2007; Madin *et al.*, 2008). De-calcification may also be exacerbated by warming ocean temperature, as is the case for coralline algae (Martin & Gattuso, 2009). To date, research on how synergies between key climate drivers affect coral reef systems is very limited (Ban *et al.*, 2014) and therefore assessments of climate change impacts usually focus on individual drivers, rather than on

cumulative effects (Cinner *et al.*, 2016). Latest IPCC projections for warm-water coral reefs globally indicate that increased water depth due to combined coral loss, reef erosion and loss of structural complexity will exacerbate the risk of flooding to reef-fringed shorelines and islands and will cause declines of reef fish populations (IPCC, 2022).

Owing to their isolation and low levels of human impacts, further studies of the Pitcairn Islands' reefs can help disentangle these synergies and provide answers to many of these scientific questions, for the benefit of this as well as other reef areas around the world.

Acknowledgements. We are very grateful for the constructive and insightful comments received from two anonymous reviewers, which helped us improve the final version of this article.

Author contributions. Susana Lincoln: Writing – Original Draft Lead Author, Review and Editing, Supervision; Benjamin Cowburn: Writing – Draft Sea Temperature Change Section; Ella L. Howes: Writing – Draft Ocean Acidification Section; Silvana Birchenough: Review – Draft Ocean Acidification Section, Review and Editing; John K. Pinnegar: Review; Stephen Dye: Writing – Draft Sea Level Rise Section; Paul Buckley: Review and Editing; Charles Sheppard: Review; Colette C. C. Wabnitz: Review; Leo X. C. Dutra: Review; Jennifer Graham: Review – Draft Sea Level Rise Section; Simeon Archer-Rand: Review; Emily Hardman: Review; Georg H. Engelhard: Review and Editing; Bryony L. Townhill: Writing – Original Draft Building Resilience to Climate Change and Knowledge Gaps and Future Research Needs, Review and Editing.

Financial support. This study was supported by the UK Government Blue Belt Programme (www.gov.uk/government/publications/the-blue-belt-programme).

Conflict of interest. The authors declare that they have no known conflict of interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- Abelson A (2020) Are we sacrificing the future of coral reefs on the altar of the 'climate change' narrative? *ICES Journal of Marine Science* 77, 40–45.
- Anthony KR, Helmstedt KJ, Bay LK, Fidelman P, Hussey KE, Lundgren P, Mead D, McLeod IM, Mumby PJ, Newlands M, Schaffelke B, Wilson KA and Hardisty PE (2020) Interventions to help coral reefs under global change – a complex decision challenge. *PLoS ONE* 15, e0236399.
- Anthony KR, Maynard JA, Diaz-Pulido G, Mumby PJ, Marshall PA, Cao L and Hoegh-Guldberg O (2011) Ocean acidification and warming will lower coral reef resilience. *Global Change Biology* 17, 1798–1808.
- Ateweberhan M, Feary DA, Keshavmurthy S, Chen A, Schleyer MH and Sheppard CRC (2013) Climate change impacts on coral reefs: synergies with local effects, possibilities for acclimation, and management implications. *Marine Pollution Bulletin* 74, 526–539.
- Australian Bureau of Meteorology and CSIRO (2014) *Climate Variability, Extremes and Change in the Western Tropical Pacific. New Science and Updated Country Reports 2014*. Melbourne: Pacific-Australia Climate Change Science and Adaptation Planning Program.
- Avagliano E, Artzner F, Kape J and Bocquet A (2016) *Regional Ecosystem Profile – Pitcairn Islands, Pacific Region*. EU Outermost Regions and Overseas Countries and Territories, BEST, Service contract 07.0307.2013/666363/SER/B2, European Commission, 60pp.
- Baird A, Madin J, Alvarez-Noriega M, Fontoura L, Kerry J, Kuo C, Precoda K, Torres-Pulliza D, Woods R, Zawada KJA and Hughes T (2018) A decline in bleaching suggests that depth can provide a refuge from global warming in most coral taxa. *Marine Ecology Progress Series* 603, 257–264. <https://doi.org/10.3354/meps12732> (Accessed online 21 January 2021).
- Baker AC, Glynn PW and Riegl B (2008) Climate change and coral reef bleaching: an ecological assessment of long-term impacts, recovery trends and future outlook. *Estuarine, Coastal and Shelf Science* 80, 435–471.
- Ban SS, Graham NAJ and Connolly SR (2014) Evidence for multiple stressor interactions and effects on coral reefs. *Global Change Biology* 20, 681–697.
- Barnett J and O'Neill SJ (2012) Islands, resettlement and adaptation. *Nature Climate Change* 2, 8–10.

- Becker M, Meysignac B, Letretel C, Llovel W, Cazenave A and Delcroix T (2012) Sea level variations at tropical Pacific islands since 1950. *Global and Planetary Change* **80–81**, 85–98.
- Bell JD, Johnson JE, Ganachaud AS, Gehrke PC, Hobday AJ, Hoegh-Guldberg O, Le Borgne R, Lehodey P, Lough JM, Pickering T, Pratchett MS and Waycott M (2011) *Vulnerability of Tropical Pacific Fisheries and Aquaculture to Climate Change: Summary for Pacific Island Countries and Territories*. New Caledonia: Secretariat of the Pacific Community, pp. 191–196.
- Belokurov A, Baskinas L, Biyo R, Clausen A, Dudley N, Guevara O, Lumanog J, Rakotondrazafy H, Ramahery V, Salao C, Stolton S and Zogib L (2016) *Climate Adaptation Methodology for Protected Areas (CAMPAs): Coastal and Marine*. Gland: WWF, 160 pp.
- Benkwitt CE, Gunn RL, Le Corre M, Carr P and Graham NAJ (2021) Rat eradication restores nutrient subsidies from seabirds across terrestrial and marine ecosystems. *Current Biology* **31**, 1–8.
- Benkwitt CE, Wilson SK and Graham NAJ (2019) Seabird nutrient subsidies alter patterns of algal abundance and fish biomass on coral reefs following a bleaching event. *Global Change Biology* **25**, 2619–2632. <https://doi.org/10.1111/gcb.14643> (Accessed online 10 January 2021).
- Bergstrom E, Silva J, Martins C and Horta P (2019) Seagrass can mitigate negative ocean acidification effects on calcifying algae. *Scientific Reports* **9**, 1–11.
- Birchenough S, Williamson P and Turley C (2017) *Future of the Sea: Ocean Acidification. Foresight – Future of the Sea. Evidence Review*. United Kingdom: Government Office for Science. Go-Science. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/645500/Ocean_Acidification_final_v3.pdf
- BirdLife International (2020) Important Bird Areas factsheet: Henderson Island. Available at <http://www.birdlife.org> (Accessed online 28 October 2020).
- Birkeland C (1989) The Faustian traits of the crown-of-thorns starfish: the structure of a fast-growing, voracious, but relatively fragile tropical predator. *American Scientist* **77**, 154–163.
- BlueBelt Report Card (2021) Climate Change Impacts on Corals in the UK Overseas Territories of BIOT and the Pitcairn Islands. In Lincoln S, Cowburn B, Howes E, Birchenough S, Pinnegar J, Dye S, Buckley P, Engelhard GE and Townhill B (eds), *UK Blue Belt Programme*, 16 pp. Available at <http://www.gov.uk/blue-belt-programme> (Accessed online 18 April 2021).
- Bruno JF, Côté IM and Toth LT (2019) Climate change, coral loss, and the curious case of the parrotfish paradigm: why don't marine protected areas improve reef resilience? *Annual Review of Marine Science* **11**, 307–334.
- Burke L, Reyter K, Spalding M and Perry A (2011) *Reefs at Risk Revisited*. Washington, DC: World Resources Institute.
- Cai W, Santoso A, Wang G, Wu L, Collin M, Lengaigne M, Power S and Timmermann A (2020) ENSO response to greenhouse forcing. In McPhaden MJ, Santoso A and Cai W (eds), *El Niño Southern Oscillation in a Changing Climate*. Available at <https://doi.org/10.1002/9781119548164.ch13> (Accessed online 2 March 2021).
- Capotondi A, Deser C, Phillips AS, Okumura Y and Larson SM (2020) ENSO and Pacific decadal variability in the community earth system model version 2. *Journal of Advances in Modelling Earth Systems* **12**, e2019MS002022.
- Carlson RR, Foo SA and Asner GP (2019) Land use impacts on coral reef health: a ridge-to-reef perspective. *Frontiers in Marine Science* **6**, 562.
- Carpenter KE, Abrar M, Aeby G, Aronson RB, Banks S, Bruckner A, Chiriboga A, Cortés J, Delbeek JC, DeVantier L, Edgar GJ, Edwards AJ, Fenner D, Guzmán HM, Hoeksema BW, Hodgson G, Johan O, Licuanan WY, Livingstone SR, Lovell ER, Moore JA, Obura DO, Ochavillo D, Polidoro BA, Precht WF, Quibilan MC, Reboton C, Richards ZT, Rogers AD, Sanciangco J, Sheppard A, Sheppard C, Smith J, Stuart S, Turak E, Veron JEN, Wallace C, Weil E and Wood E (2008) One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science (New York, N.Y.)* **321**, 560–563.
- Cazenave A and Remy F (2011) Sea level and climate: measurements and causes of changes. *WIREs Climate Change* **2**, 647–662.
- Cerutti JMB, Burt AJ, Haupt P, Bunbury N, Mumby P and Schaeppman-Strub G (2020) Impacts of the 2014–2017 global bleaching event on a protected remote atoll in the Western Indian Ocean. *Coral Reefs* **39**, 15–26.
- Chan NC and Connolly SR (2013) Sensitivity of coral calcification to ocean acidification: a meta-analysis. *Global Change Biology* **19**, 282–290.
- Chand SS, Dowdy A, Bell S and Tory K (2020) A review of South Pacific tropical cyclones: impacts of natural climate variability and climate change. In Kumar L (ed.), *Climate Change and Impacts in the Pacific*. Cham: Springer, pp. 251–273.
- Christensen JH, Kumar KK, Aldrian E, An SI, Cavalcanti IFA, de Castro M, Dong W, Goswami P, Hall A, Kanyanga JK, Kitoh A, Kossin J, Lau NC, Renwick J, Stephenson DB, Xie SP and Zhou T (2013) Climate phenomena and their relevance for future regional climate change. Supplementary material. In Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 1217–1308.
- Church JA, Clark PU, Cazenave A, Gregory JM, Jevrejeva S, Levermann A, Merrifield MA, Milne GA, Nerem RS, Nunn PD, Payne AJ, Pfeffer WT, Stammer D and Unnikrishnan AS (2013) Sea level change. In Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds), *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 1137–1216.
- Cinner JE, Pratchett MS, Graham NAJ, Messmer V, Fuentes MMPB, Ainsworth T, Ban N, Bay LK, Blythe J, Dissard D, Dunn S, Evans L, Fabinyi M, Fidelman P, Figueiredo J, Frisch AJ, Fulton CJ, Hicks CC, Lukoschek V, Mallela J, Moya A, Penin L, Rummer JL, Walker S and Williamson DH (2016) A framework for understanding climate change impacts on coral reef social-ecological systems. *Regional Environmental Change* **16**, 1133–1146.
- Coghlan AR, White R, Dawson TP, Irving RA, Zeller D and Palomares ML (2017) Reconstructed marine fisheries catches at a remote island group: Pitcairn Islands (1950–2014). *Frontiers in Marine Science* **4**, 320.
- Collins M, Sutherland M, Bouwer L, Cheong S-M, Frölicher T, Jacot Des Combes H, Roxy MK, Losada I, McInnes K, Ratter B, Rivera-Arriaga E, Susanto RD, Swingedouw D and Tibig L (2019) Extremes, abrupt changes and managing risk. In Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M, Poloczanska E, Mintenbeck K, Alegria A, Nicolai M, Okem A, Petzold J, Rama B and Weyer NM (eds), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Available at <https://www.ipcc.ch/srocc/cite-report/> (Accessed online 15 March 2021).
- Cornwall W (2019) Researchers embrace a radical idea: engineering coral to cope with climate change. *Science. Climate News* (21 Mar 2019). <https://doi.org/10.1126/science.aax4091> (Accessed online 2 March 2021).
- Couce E, Ridgwell A and Hendy EJ (2013) Future habitat suitability for coral reef ecosystems under global warming and ocean acidification. *Global Change Biology* **19**, 3592–3606.
- Cowburn B, Moritz C, Grimsditch G and Solandt J (2019) Evidence of coral bleaching avoidance, resistance and recovery in the Maldives during the 2016 mass-bleaching event. *Marine Ecology Progress Series* **626**, 53–67.
- Crabbe MJC (2019) Adapting to extreme environments: can coral reefs adapt to climate change? *Emerging Topics in Life Sciences* **3**, 183–195.
- CSIRO, Australian Bureau of Meteorology and SPREP (2015) *Climate in the Pacific: A Regional Summary of New Science and Management Tools*. Melbourne: Commonwealth Scientific and Industrial Research Organisation.
- Cyronak T, Andersson AJ, D'Angelo S, Bresnahan P, Davidson C, Griffin A, Kindeberg T, Pennise J, Takeshita Y and White M (2018) Short-term spatial and temporal carbonate chemistry variability in two contrasting seagrass meadows: implications for pH buffering capacities. *Estuaries and Coasts* **41**, 1282–1296.
- Dance A (2019) These corals could survive climate change – and help save the world's reefs. *Nature* **575**, 580–582.
- Dawson TP and Irving RA (2020) Chapter 14 - Developing a fisheries management plan for the Pitcairn Islands Marine Reserve. In Humphreys J and Clark RWE (eds), *Marine Protected Areas, Science Policy and Management*. United Kingdom: Elsevier, pp. 271–283.
- Dawson TP and Zhang X (2020) Climate impacts on the coral reefs of the Pitcairn Islands Marine Reserve in 2016. *MASTS 10th Annual Science Meeting 2020 'MASTS – A decade of innovation', 5–9th October*. Marine Alliance for Science and Technology for Scotland.

- Diamond HJ, Lorrey AM and Renwick JA (2013) A southwest Pacific tropical cyclone climatology and linkages to the El Niño-Southern Oscillation. *Journal of Climate* **26**, 3–25.
- Doney SC, Busch DS, Cooley SR and Kroeker KJ (2020) The impacts of ocean acidification on marine ecosystems and reliant human communities. *Annual Review of Environment and Resources* **45**, 83–112.
- Doo SS, Kealoha A, Andersson A, Cohen AL, Hicks TL, Johnson ZI, Long MH, Mcelhany P, Mollica N, Shamberger KEF, Silbiger NJ, Takeshita Y, Busch DS and Browman H (2020) The challenges of detecting and attributing ocean acidification impacts on marine ecosystems. *ICES Journal of Marine Science* **77**, 2144–2422. doi: 10.1093/icesjms/fsaa094
- Doropoulos C, Vons F, Elzinga J, ter Hofstede R, Salee K, van Koningsveld M and Babcock RC (2019) Testing industrial-scale coral restoration techniques: harvesting and culturing wild coral-spawn slicks. *Frontiers in Marine Science* **6**, 1–14.
- Duffy H (2014) *Fisheries Resources of Pitcairn Island: Assessment and Management Challenges*, MSc Thesis in Conservation Science. London, United Kingdom: Imperial College London.
- Duffy H, Letessier TB, Koldewey HJ, Dawson TP and Irving RA (2021): Ensuring the sustainability of coastal small-scale fisheries at Pitcairn Island (South Pacific) within a large scale no-take MPA. *Frontiers in Marine Science* **8**, 528.
- Dutheil C, Lengaigne M, Bador M, Vialard J, Lefèvre J, Jourdain NC, Jullien S, Peltier A, Sultan B and Menkès C (2020) Impact of projected sea surface temperature biases on tropical cyclones projections in the South Pacific. *Scientific Reports* **10**, 4838.
- Dutra LXC, Haywood MDE, Singh S, Ferreira M, Johnson JE, Veitayaki J, Kininmonth S, Morris CW and Piovano S (2021) Synergies between local and climate-driven impacts on coral reefs in the Tropical Pacific: a review of issues and adaptation opportunities. *Marine Pollution Bulletin* **164**, 111922.
- Dutra LXC, Haywood MDE, Singh SS, Piovano S, Ferreira M, Johnson JE, Veitayaki J, Kininmonth S and Morris CW (2018) Impacts of climate change on corals relevant to the Pacific Islands. *Pacific Marine Climate Change Report Card: Science Review* **2018**, 132–158.
- Enochs IC, Manzello DP, Donham EM, Kolodziej G, Okano R, Johnston L, Young C, Iguel J, Edwards CB, Fox MD, Valentino L, Johnson S, Benavente D, Clark SJ, Carlton R, Burton T, Eynaud Y and Price NN (2015) Shift from coral to macroalgae dominance on a volcanically acidified reef. *Nature Climate Change* **5**, 1083–1088.
- Enochs IC, Manzello DP, Kolodziej G, Noonan SH, Valentino L and Fabricius KE (2016) Enhanced macroboring and depressed calcification drive net dissolution at high-CO₂ coral reefs. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20161742.
- Fabricius KE, Langdon C, Uthicke S, Humphrey C, Noonan S, De'ath G, Okazaki R, Muehlehner N, Glas MS and Lough JM (2011) Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. *Nature Climate Change* **1**, 165–169.
- Fabricius KE, Neill C, Van Ooijen E, Smith JN and Tilbrook B (2020) Progressive seawater acidification on the Great Barrier Reef continental shelf. *Scientific Reports* **10**, 1–15.
- Fang SW and Yu JY (2020) A control of ENSO transition complexity by tropical Pacific mean SSTs through tropical-subtropical interaction. *Geophysical Research Letters* **47**, e2020GL087933.
- Feely RA, Doney SC and Cooley SR (2009) Ocean acidification: present conditions and future changes in a high-CO₂ world. *Oceanography* **22**, 36–47.
- Fiedler P and Lavin M (2017) Oceanographic conditions of the eastern tropical Pacific. In Glynn PW, Manzello D and Enochs IC (eds), *Coral Reefs of the Eastern Tropical Pacific: Persistence and Loss in a Dynamic Environment*. Coral Reefs of the World 8. Dordrecht: Springer Science, 666 pp. Available at https://doi.org/10.1007/978-94-017-7499-4_3 (Accessed online 21 January 2021).
- Field CB, Barros VR, Mastrandrea MD, Mach KJ, Abdrabo M-K, Adger N, Anokhin YA, Anisimov OA, Arent DJ and Barnett J (2014) *Summary for Policymakers. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom: Cambridge University Press.
- Fisher R, O'Leary RA, Low-Choy S, Mengersen K, Knowlton N, Brainard RE and Caley MJ (2015) Species richness on coral reefs and the pursuit of convergent global estimates. *Current Biology* **25**, 500–505.
- Fletcher CH, Bochicchio C, Conger CL, Engels MS, Feirstein EJ, Frazer N, Glenn CR, Grigg RW, Grossman EE, Harney JN, Isoun E, Murray-Wallace CV, Rooney JJ, Rubin KH, Sherman CE and Vitousek S (2008) Geology of Hawaii reefs. In Riegl B and Dodge R (eds), *Coral Reefs of the USA*. Dordrecht: Springer, pp. 435–487.
- Ford MR, Kench PS, Owen SD and Hua Q (2020) Active sediment generation on coral reef flats contributes to recent reef island expansion. *Geophysical Research Letters* **47**, e2020GL088752.
- Forrest AK and Hindell M (2018) Ingestion of plastic by fish destined for human consumption in remote South Pacific Islands. *Australian Journal of Maritime & Ocean Affairs* **10**, 81–97.
- Friedlander AM, Caselle JE, Ballesteros E, Brown EK, Turchik A and Sala E (2014) The real bounty: marine biodiversity in the Pitcairn Islands. *PLoS ONE* **9**, e100142.
- Galloway SB, Bruckner AW and Woodley CM (2009) *Coral Health and Disease in the Pacific: Vision for Action*. NOAA Technical Memorandum NOS NCCOS 97 and CRCP 7. Silver Spring, MD: National Oceanic and Atmospheric Administration, p. 314. Available at www.coris.noaa.gov/activities/cdh_vision/pdfs/cdh_vision_full.pdf (Accessed online 15 January 2021).
- Giglio V, Ternes M, Mendes T, Cordeiro C and Ferreira C (2017) Anchoring damages to benthic organisms in a subtropical scuba dive hotspot. *Journal of Coastal Conservation* **21**, 311–316.
- Gilmour JP, Smith LD, Heyward AJ, Baird AH and Pratchett MS (2013) Recovery of an isolated coral reef system following severe disturbance. *Science (New York, N.Y.)* **340**, 69–71.
- Glynn PW (1984) Widespread coral mortality and the 1982–83 El Niño warming event. *Environment Conservation* **11**, 133–146.
- Glynn PW (1993) Coral reef bleaching: ecological perspectives. *Coral Reefs* **12**, 1–17.
- GOA-ON (2020) *Global Ocean Acidification Observing Network*. Available at <http://goa-on.org> (Accessed online 23 March 2021).
- The Government of the Pitcairn Islands (2017) *Revised Laws of Pitcairn, Henderson, Ducie and Oeno Islands, 2017*, Revised Edition. Volume 2. Available at <http://government.pn/Laws/Cap%2048%20-%20Pitcairn%20Islands%20Marine%20Protected%20Area%202017%20Rev%20Ed.pdf#:~:text=The%20following%20areas%20are%20established%20as%20Coastal%20Conservation,mile%20reef%20as%20described%20in%20Marine%20Conservation%20Regulations>.
- Graham NAJ, McClanahan TR, MacNeil MA, Wilson SK, Cinner JE, Huchery C and Holmes TH (2017) Human disruption of coral reef trophic structure. *Current Biology* **27**, 231–236.
- Graham NAJ, Wilson SK, Carr P, Hoey AS, Jennings S and MacNeil MA (2018) Seabirds enhance coral reef productivity and functioning in the absence of invasive rats. *Nature* **559**, 250–253.
- Groose MR, Narsey S, Delage FP, Dowdy AJ, Bador M, Boschat G, Chung C, Kajtar JB, Rauniyar S, Freund MB, Lyu K, Rashid H, Zhang X, Wales S, Trenham C, Holbrook NJ, Cowan T, Alexander L, Arblaster JM and Power S (2020) Insights from CMIP6 for Australia's future climate. *Earth's Future* **8**, e2019EF001469.
- Harvey BJ, Nash KL, Blanchard JL and Edwards DP (2018) Ecosystem-based management of coral reefs under climate change. *Ecology and Evolution* **8**, 6354–6368.
- Hays GC, Koldewey HJ, Andrzejczak S, Attrill MJ, Barley S, Bayley DTI, Benkowitz CE, Block B, Schallert RJ, Carlisle AB, Carr P, Chapple TK, Collins C, Diaz C, Dunn N, Dunbar RB, Eager DS, Engel J, Embling CB, Esteban N, Ferretti F, Foster NL, Freeman R, Gollock M, Graham NAJ, Harris JL, Head CEI, Hosegood P, Howell KL, Hussey NE, Jacoby DMP, Jones R, Pilly SS, Lange ID, Letessier TB, Levy E, Lindhart M, McDevitt-Irwin JM, Meekam M, Meeuwig JJ, Micheli F, Mogg AOM, Mortimer JA, Mucciarone DA, Nicoll MA, Nuno A, Perry CT, Preston SG, Rattray AJ, Robinson E, Roche RC, Schiele M, Sheehan EV, Sheppard A, Sheppard C, Smith AL, Soule B, Spalding M, Stevens GMW, Steyaert M, Stifel S, Taylor BM, Tickler D, Trevail AM, Trueba P, Turner J, Votier S, Wilson B, Williams GJ, Williamson BJ, Williamson MJ, Wood H and Curnick DJ (2020) A review of a decade of lessons from one of the world's largest MPAs: conservation gains and key challenges. *Marine Biology* **167**, 159.
- Head CEI, Bayley DTI, Rowlands G, Roche RC, Tickler DM, Rogers AD, Koldewey H, Turner JR and Andradi-Brown DA (2019) Coral bleaching impacts from back-to-back 2015–2016 thermal anomalies in the remote central Indian Ocean. *Coral Reefs* **38**, 605–618.
- Heron S, Maynard J, Van Hooidonk R and Eaking M (2016) Warming trends and bleaching stress of the world's coral reefs 1985–2012. *Scientific Reports* **6**, 38402.

- Hoarau K, Chalonge L, Pirard F and Peyrusaubes D (2018) Extreme tropical cyclone activities in the Southern Pacific Ocean. *International Journal of Climatology* **38**, 1409–1420.
- Hoegh-Guldberg O and Bruno JF (2010) The impact of climate change on the world's marine ecosystems. *Science (New York, N.Y.)* **328**, 1523–1528.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A and Hatzioles ME (2007) Coral reefs under rapid climate change and ocean acidification. *Science (New York, N.Y.)* **1737**, 1737–1743.
- Holland G and Bruyère CL (2014) Recent intense hurricane response to global climate change. *Climate Dynamics* **42**, 617–627.
- Hughes TP, Anderson KD, Connolly SR, Heron SF, Kerry JT, Lough JM, Baird AH, Baum JK, Berumen ML, Bridge TC and Claar DC (2018) Spatial and temporal patterns of mass bleaching of corals in the Anthropocene. *Science (New York, N.Y.)* **359**, 80–83.
- Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson JB, Kleypas J, Lough JM, Marshall P, Nystrom M, Palumbi SR, Pandolfi JM, Rosen B and Roughgarden J (2003) Climate change, human impacts, and the resilience of coral reefs. *Science (New York, N.Y.)* **301**, 929–933.
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, Babcock RC, Beger M, Bellwood DR, Berkelmans R, Bridge TC, Butler IR, Byrne M, Cantin NE, Comeau S, Connolly SR, Cumming GS, Dalton SJ, Diaz-Pulido G, Eakin CM, Figueira WF, Gilmour JP, Harrison HB, Heron SF, Hoey AS, Hobbs JPA, Hoogenboom MO, Kennedy EV, Kuo CY, Lough JM, Lowe RJ, Liu G, McCulloch MT, Malcolm HA, McWilliam MJ, Pandolfi JM, Pears RJ, Pratchett MS, Schoepf V, Simpson T, Skirving WJ, Sommer B, Torda G, Wachenfeld DR, Willis BL and Wilson SK (2017) Global warming and recurrent mass bleaching of corals. *Nature* **543**, 373–377.
- Hurd CL, Lenton A, Tilbrook B and Boyd PW (2018) Current understanding and challenges for oceans in a higher-CO₂ world. *Nature Climate Change* **8**, 686–694.
- IPCC (2013) Climate change 2013: the physical science basis. In Stocker TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V and Midgley PM (eds), *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 1–1535.
- IPCC (2014a) Climate change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects. In Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR and White LL (eds), *Working Group II Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 688.
- IPCC (2014b) Climate change 2014: synthesis report. In Pachauri RK and Meyer LA (eds), *Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, vol. 218. Geneva: IPCC, 151 pp.
- IPCC (2019a) Chapter 4: sea level rise and implications for low lying islands, coasts and communities. In Pörtner DCRH-O, Masson-Delmotte V, Zhai P, Tignor M, Poloczanska E, Mintenbeck K, Nicolai M, Okem A, Petzold J, Rama B and Weyer N (eds), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Cambridge: Cambridge University Press, pp. 321–446.
- IPCC (2019b) Summary for policymakers. In Pörtner H-O, Roberts DC, Masson-Delmotte V, Zhai P, Tignor M, Poloczanska E, Mintenbeck K, Nicolai M, Okem A, Petzold J, Rama B and Weyer N (eds), *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*. Cambridge: Cambridge University Press, pp. 3–35.
- IPCC (2021) Climate change 2021: the physical science basis. In Masson-Delmotte VP, Zhai A, Pirani SL, Connors C, Péan S, Berger N, Caud Y, Chen L, Goldfarb MI, Gomis M, Huang K, Leitzell E, Lonnoy JBR, Matthews TK, Maycock T, Waterfield O, Yelekçi RY and Zhou B (eds), *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 2391.
- IPCC (2022) Climate change 2022: impacts, adaptation and vulnerability. In Pörtner H-O, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Lösschke S, Möller V, Okem A and Rama B (eds), *Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 3056 pp. doi: 10.1017/9781009325844.
- Irving RA (1995) Near-shore bathymetry and reef biotopes of Henderson Island, Pitcairn Group. *Biological Journal of the Linnean Society* **56**, 309–324.
- Irving R and Dawson TP (2012) *The Marine Environment of the Pitcairn Islands. A Report to Global Ocean Legacy, a Project of the Pew Environment Group*. Available at <https://www.pewtrusts.org/-/media/legacy/uploadedfiles/peg/publications/report/dupreport11lopdf.pdf>.
- Irving R and Dawson TP (2013) Coral reefs of the Pitcairn Islands. In Sheppard CRC (ed.), *Coral Reefs of the United Kingdom Overseas Territories*. Berlin: Springer-Verlag, pp. 253–270.
- Irving RA, Dawson TP and Christian M (2019) Chapter 34 – The Pitcairn Islands. In Sheppard C (ed.), *World Seas: An Environmental Evaluation*, 2nd Edition. London, United Kingdom: Academic Press, pp. 743–764.
- Irving RA, Dawson TP and Wovor D (2017) An amphidromic prawn, *Macrobrachium latimanus* (von Martens, 1868) (Decapoda: Palaemonidae), discovered on Pitcairn, a remote island in the southeastern Pacific Ocean. *Journal of Crustacean Biology* **37**, 503–506.
- Johnson J, Bell J and Gupta AS (2015) *Pacific Islands Ocean Acidification Vulnerability Assessment*. Apia, Samoa: SPREP, 40 pp.
- Jones LA, Mannion PD, Farnsworth A, Valdes PJ, Kelland S and Allison PA (2019) Coupling of palaeontological and neontological reef coral data improves forecasts of biodiversity responses under global climatic change. *Royal Society Open Science* **6**, 182111.
- Kench PS, Ford MR and Owen SD (2018a) Patterns of island change and persistence offer alternate adaptation pathways for atoll nations. *Nature Communications* **9**, 605. <https://doi.org/10.1038/s41467-018-02954-1>
- Kench PS, McLean RF, Owen SD, Tuck M and Ford MR (2018b) Storm-deposited coral blocks: a mechanism of island genesis, Tutaga island, Funafuti atoll, Tuvalu. *Geology* **46**, 915–918.
- Kench PS, Thompson D, Ford MR, Ogawa H and McLean RF (2015) Coral islands defy sea level rise over the past century: records from a Central Pacific atoll. *Geology* **43**, 515–518.
- Kleypas JA, McManus JW and Meñez LAB (2010) Environmental limits to coral reef development: where do we draw the line? *American Zoologist* **39**, 146–159.
- Kubota K, Yokoyama Y, Ishikawa T, Suzuki A and Ishii M (2017) Rapid decline in pH of coral calcification fluid due to incorporation of anthropogenic CO₂. *Scientific Reports* **7**, 7694.
- Lafale PF, Diamond HJ and Anderson CL (2018) Impacts of climate change on extreme events relevant to the Pacific Islands. *Pacific Marine Climate Change Report Card: Science Review* **2018**, 50–73.
- Lamb JB, Willis BL, Fiorenza EA, Couch CS, Howard R, Rader DN, True JD, Kelly LA, Ahmad A, Jompa J and Harvell CD (2018) Plastic waste associated with disease on coral reefs. *Science (New York, N.Y.)* **359**, 460–462.
- Langdon C and Atkinson M (2005) Effect of elevated pCO₂ on photosynthesis and calcification of corals and interactions with seasonal change in temperature/irradiance and nutrient enrichment. *Journal of Geophysical Research: Oceans* **110**, C09S07.
- Lauvsek SK, Gruber N, Landschützer P, Olsen A and Tjiputra J (2015) Trends and drivers in global surface ocean pH over the past 3 decades. *Biogeosciences (Online)* **12**, 1285–1298.
- Lavers JL and Bond AL (2017) Significant anthropogenic debris on remote island. *Proceedings of the National Academy of Sciences USA* **114**, 6052–6055.
- Lavers JL, Rivers-Auty J and Bond AL (2021) Plastic debris increases circadian temperature extremes in beach sediments. *Journal of Hazardous Materials* **416**, 126140.
- Lenton A, Matear RL and Mongin M (2018) Impacts of climate change on ocean acidification relevant to the Pacific Islands. *Pacific Marine Climate Change Report Card: Science Review* **2018**, 31–42.
- Lewis N, Day JC, Wilhelm A, Wagner D, Gaymer C, Parks J, Friedlander A, White S, Sheppard C, Spalding M, San Martin G, Skeat A, Taii S, Teroroko T and Evans J (2017) *Large-Scale Marine Protected Areas: Guidelines for Design and Management. Best Practice Protected Area Guidelines Series, No. 26*. Gland: IUCN, xxviii+120 pp.
- Lincoln S, Cowburn B, Howes E, Birchenough SNR, Pinnegar J, Dye S, Buckley P, Engelhard GH and Townhill BL (2021) *Climate Change Impacts on Corals in the UK Overseas Territories of BIOT and the Pitcairn Islands. Scientific Report*. UK Blue Belt Programme, 83 pp. Available at <http://www.gov.uk/blue-belt-programme>.

- Liu PJ, Ang SJ, Mayfield AB and Lin HJ (2020) Influence of the seagrass *Thalassia hemprichii* on coral reef mesocosms exposed to ocean acidification and experimentally elevated temperatures. *Science of the Total Environment* **700**, 13464.
- Loya Y, Sakai K, Yamazato K, Nakano Y, Sambali H and van Woesik R (2001) Coral bleaching: the winners and the losers. *Ecology Letters* **4**, 122–131.
- MacNeil A, Mellin C, Matthews S, Wolff N, McClanahan T, Devlin M, Drovandi C, Mengersen K and Graham N (2019) Water quality mediates resilience on the Great Barrier Reef. *Nature Ecology & Evolution* **3**, 620–627. <https://doi.org/10.1038/s41559-019-0832-3>
- Madin JS, O'Donnell MJ and Connolly SR (2008) Climate-mediated mechanical changes to post-disturbance coral assemblages. *Biology Letters* **4**, 490–493.
- Martin S and Gattuso J-P (2009) Response of Mediterranean coralline algae to ocean acidification and elevated temperature. *Global Change Biology* **15**, 2089–2099.
- Masselink G, Beetham E and Kench P (2020) Coral reef islands can accrete vertically in response to sea level rise. *Science Advances* **6**, eaay3656.
- McClanahan TR, Cinner JE, Maina J, Graham NAJ, Daw TM, Stead SM, Wamukota A, Brown K, Ateweberhan M, Venus V and Polunin NVC (2008) Conservation action in a changing climate. *Conservation Letters* **1**, 53–59.
- McLean R and Kench P (2015) Destruction or persistence of coral atoll islands in the face of 20th and 21st century sea level rise? *Wiley Interdisciplinary Reviews: Climate Change* **6**, 445–463.
- McLeod E, Anthony KRN, Mumby PJ, Maynard J, Beeden R, Graham NAJ, Heron SF, Hoegh-Guldberg O, Jupiter S, MacGowan P, Mangubhai S, Marshall N, Marshall PA, McClanahan TR, McLeod K, Nyström M, Obura D, Parker B, Possingham HP, Salm RV and Tamelander J (2019) The future of resilience-based management in coral reef ecosystems. *Journal of Environmental Management* **233**, 291–301.
- McPhaden MJ, Zebiak SE and Glantz MH (2006) ENSO as an integrating concept in earth. *Science (New York, N.Y.)* **314**, 1740–1745.
- Meyssignac B, Becker M, Llovel W and Cazenave A (2012) An assessment of two-dimensional past sea level reconstructions over 1950–2009 based on tide-gauge data and different input sea level grids. *Surveys in Geophysics* **33**, 945–972.
- Moberg F and Folke C (1999) Ecological goods and services of coral reef ecosystems. *Ecological Economics* **29**, 215–233.
- Mollica NR, Guo W, Cohen AL, Huang KF, Foster GL, Donald HK and Solow AR (2018) Ocean acidification affects coral growth by reducing skeletal density. *Proceedings of the National Academy of Sciences USA* **115**, 1754–1759.
- Moritz C, Vii J, Long WL, Tamelander J, Thomassin A and Planes S (eds) (2018) *Status and Trends of Coral Reefs of the Pacific*. Global Coral Reef Monitoring Network.
- Muir PR, Marshall PA, Abdulla A and Aguirre JD (2017) Species identity and depth predict bleaching severity in reef-building corals: shall the deep inherit the reef? *Proceedings of the Royal Society B: Biological Sciences* **284**, 20171551.
- Mumby PJ (2009) Herbivory versus corallivory: are parrotfish good or bad for Caribbean coral reefs? *Coral Reefs* **28**, 683–690.
- Mumby PJ, Hastings A and Edwards HJ (2007) Thresholds and the resilience of Caribbean coral reefs. *Nature Letters* **450**, 98–101.
- Mumby PJ, Steneck RS, Adjeroud M and Arnold SN (2016) High resilience masks underlying sensitivity to algal phase shifts of Pacific coral reefs. *Oikos* **125**, 644–655.
- Nakamura T and Van Woesik R (2001) Water-flow rates and passive diffusion partially explain differential survival of corals during the 1998 bleaching event. *Marine Ecology Progress Series* **212**, 301–304.
- NOAA Climate Change Portal (2020) CIMP5. Available at <https://psl.noaa.gov/ipcc/ocn/> (Accessed online 7 October 2020).
- NOAA Coral Reef Conservation Program (2014) *Climate Change Strategy and Implementation Plan, 2014–2016*. Silver Spring, MD, 28 pp. Available at https://www.ncei.noaa.gov/data/oceans/coris/library/NOAA/CRCP/project/851/CRCP_CCSIP_Final.pdf.
- NOAA Coral Reef Watch (2018) *NOAA Coral Reef Watch Version 3.1 Daily Global 5-km Satellite Coral Bleaching Degree Heating Week Product*. College Park, MD: NOAA Coral Reef Watch. Available at <https://coralreefwatch.noaa.gov/product/5km/index.php> (Accessed online 1 September 2018).
- Nurse L, McLean R, Agard J and Briguglio L (2014) Small islands. In Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR and White LL (eds), *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 1613–1654.
- Nyström M, Graham NAJ, Lokrantz J and Norström A (2008) Capturing the cornerstones of coral reef resilience: linking theory to practice. *Coral Reefs* **27**, 795–809.
- Obura D and Grimsditch G (2009) *Coral Reefs, Climate Change and Resilience – An Agenda for Action from the IUCN World Conservation Congress*. October 6–9, 2008, 44 pp. Available at <https://portals.iucn.org/library/sites/library/files/documents/2009-022.pdf>.
- Pacella SR, Brown CA, Waldbusser GG, Labiosa RG and Hales B (2018) Seagrass habitat metabolism increases short-term extremes and long-term offset of CO₂ under future ocean acidification. *Proceedings of the National Academy of Sciences USA* **115**, 3870–3875.
- Pandolfi JM, Connolly SR, Marshall DJ and Cohen AL (2011) Projecting coral reef futures under global warming and ocean acidification. *Science (New York, N.Y.)* **333**, 418–422.
- Perez-Correa J, Carr P, Meeuwij JJ, Koldewey HJ and Letessier TB (2020) Climate oscillation and the invasion of alien species influence the oceanic distribution of seabirds. *Ecology and Evolution* **10**, 9339–9357.
- Pitcairn Government (2017) Pitcairn, Henderson, Ducie and Oeno Islands: Marine Protected Areas. In *Revised Laws of Pitcairn, Henderson, Ducie and Oeno Islands*, 2017 Rev. Ed. – Volume 2 (government.pn) (Accessed online 14 April 2021).
- Pitcairn Miscellany (2020) *The Pitcairn Miscellany, January 2020*. Available at <http://www.miscellany.pn> (Accessed online 16 April 2021).
- Pratchett MS, McCowan D, Maynard JA and Heron SF (2013) Changes in bleaching susceptibility among corals subject to ocean warming and recurrent bleaching in Moorea, French Polynesia. *PLoS ONE* **8**, 1–10.
- PSMSL (2020) *Tide Gauge Data, Permanent Service for Mean Sea Level*. Available at <http://www.psmsl.org/data/obtaining> (Accessed online 23 October 2020).
- Readman JW, Deluna F, Ebinghaus R, Guzman A, Price A, Readman E, Sheppard A, Sleight V, Sturm R, Thompson R, Tonkin A, Wolschke H, Wright R and Sheppard C (2013) Contaminants, pollution and potential anthropogenic impacts in Chagos/BIOF. In Sheppard CRC (ed.), *Coral Reefs of the United Kingdom Overseas Territories, Coral Reefs of the World 4*, 283. Dordrecht: Springer Science+Business Media. Available at https://doi.org/10.1007/978-94-007-5965-7_21.
- Rehder HA and Randall JE (1975) Atoll research bulletin No. 183 Ducie Atoll: its history, physiography and biota. *Atoll Research Bulletin* **183**, 1–57.
- Robinson J, New AL, Popova EE, Srokosz MA and Yool A (2017) Far-field connectivity of the UK's four largest marine protected areas: four of a kind? *Earth's Future* **5**, 475–494.
- Roche RC, Williams GJ and Turner JR (2018) Towards developing a mechanistic understanding of coral reef resilience to thermal stress across multiple scales. *Current Climate Change Reports* **4**, 51–64.
- Roff G, Doropoulos C, Rogers A, Bozec Y-M, Krueck NC, Aurellado E, Priest M, Birrell C and Mumby PJ (2016) The ecological role of sharks on coral reefs. *Trends in Ecology & Evolution* **31**, 395–407.
- Romero-Torres M, Treml EA, Acosta A and Paz-García DA (2018) The eastern tropical Pacific coral population connectivity and the role of the eastern Pacific barrier. *Scientific Reports* **8**, 1–13.
- Rooney J, Fletcher C, Grossman E, Engels M and Field M (2004) El Niño influence on Holocene reef accretion in Hawai'i. *Pacific Science* **58**, 305–324.
- Ryan PG (2020) Using photographs to record plastic in seabird nests. *Marine Pollution Bulletin* **156**, 111262.
- Ryan PG and Schofield A (2020) Low densities of macroplastic debris in the Pitcairn Islands Marine Reserve. *Marine Pollution Bulletin* **157**, 111373.
- Santoso A, McGregor S, Jin FF, Cai W, England MH, An SI, McPhaden MJ and Guilyardi E (2013) Late-twentieth-century emergence of the El Niño propagation asymmetry and future projections. *Nature* **504**, 126–130.
- Sea Around Us (2016) *Catches by Taxon in the Waters of Pitcairn (UK)*. Sea Around Us. Fisheries, Ecosystems and Biodiversity (Accessed online 17 March 2021).
- Sellar AA, Jones CG, Mulcahy JP, Tang Y, Yool A, Wiltshire A, O'Connor FM, Stringer M, Hill R, Palmieri J, Woodward S, de Mora L, Kuhlbrodt T, Rumbold ST, Kelley DI, Ellis R, Johnson CE, Walton J, Abraham NL, Andrews MB, Andrews T, Archibald AT, Berthou S, Burke E, Blockley E, Carslaw K, Dalvi M, Edwards J, Folberth GA, Gedney N, Griffiths PT,

- Harper AB, Hendry MA, Hewitt AJ, Johnson B, Jones A, Jones CD, Keeble J, Liddicoat S, Morgenstern O, Parker RJ, Predoi V, Robertson E, Siahaan A, Smith RS, Swaminathan R, Woodhouse MT, Zeng G and Zerroukat M (2019) UKESM1: description and evaluation of the UK earth system model. *Journal of Advances in Modeling Earth Systems* **11**, 4513–4558.
- Seneviratne SI, Nicholls N, Easterling D, Goodess CM, Kanae S, Kossin J, Luo Y, Marengo J, McInnes K, Rahimi M, Reichstein M, Sorteberg A, Vera C and Zhang X (2012) Changes in climate extremes and their impacts on the natural physical environment. In Field CB, Barros V, Stocker TF, Qin D, Dokken DJ, Ebi KL, Mastrandrea MD, Mach KJ, Plattner G-K, Allen SK, Tignor M and Midgley PM (eds), *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, pp. 109–230.
- Sheppard CRC, Harris A and Sheppard ALS (2008) Archipelago-wide coral recovery patterns since 1998 in the Chagos Archipelago, central Indian Ocean. *Marine Ecology Progress Series* **362**, 109–117.
- Sheppard C and Sheppard A (2019) Chapter 11 – British Indian Ocean Territory (Chagos Archipelago). In Sheppard C (ed.), *World Seas: An Environmental Evaluation*, 2nd Edn. Academic Press, pp. 237–252. Available at <https://doi.org/10.1016/B978-0-08-100853-9.00015-4>.
- Sheppard C, Sheppard A and Fenner D (2020) Coral mass mortalities in the Chagos Archipelago over 40 years: regional species and assemblage extinctions and indications of positive feedbacks. *Marine Pollution Bulletin* **154**, 111075.
- Smith TB, Glynn PW, Maté JL, Toth LT and Gyory J (2014) A depth refugium from catastrophic coral bleaching prevents regional extinction. *Ecology* **95**, 1663–1673.
- Smith TB, Gyory J, Brandt ME, Miller WJ, Jossart J and Nemeth RS (2016) Caribbean mesophotic coral ecosystems are unlikely climate change refugia. *Global Change Biology* **22**, 2756–2765.
- Spencer T (1995) The Pitcairn Islands, South Pacific Ocean: plate tectonic and climatic contexts. *Biological Journal of the Linnean Society* **56**, 13–42.
- Steneck RS, Arnold SN, Boenish R, de León R, Mumby PJ, Rasher DB and Wilson MW (2019) Managing recovery resilience in coral reefs against climate-induced bleaching and hurricanes: a 15 year case study from Bonaire, Dutch Caribbean. *Frontiers in Marine Science* **6**, 265.
- Stephens SA and Ramsay DL (2014) Extreme cyclone wave climate in the southwest Pacific Ocean: influence of the El Niño southern oscillation and projected climate change. *Global and Planetary Change* **123**, 13–26.
- Storz CD, Gingerich SB, van Dongeren A, Cheriton OM, Swarzenski PW, Quataert E, Voss CI, Field DW, Annamalai H, Piniak GA and McCall R (2018) Most atolls will be uninhabitable by the mid-21st century because of sea-level rise exacerbating wave-driven flooding. *Science Advances* **4**, eaap9741.
- Sully S, Burkepile DE, Donovan MK, Hodgson G and Van Woesik R (2019) A global analysis of coral bleaching over the past two decades. *Nature Communications* **10**, 1264. <https://doi.org/10.1038/s41467-019-09238-2>.
- Sutton PJH (2018) Impacts of climate change on Sea temperature relevant to the Pacific Islands. *Pacific Marine Climate Change Report Card: Science Review* **2018**, 20–30.
- Sutton AJ, Feely RA, Sabine CL, McPhaden MJ, Takahashi T, Chavez FP, Friederich GE and Mathis JT (2014) Natural variability and anthropogenic change in equatorial Pacific surface ocean pCO₂ and pH. *Global Biogeochemical Cycles* **28**, 131–145.
- Swearer SE, Caselle JE, Lea DW and Warner RR (1999) Larval retention and recruitment in an island population of a coral-reef fish. *Nature* **402**, 799–802.
- Tajima Y, Shoji T and Taniguchi K (2020) Study on probabilistic inundation hazard along the coast of South Pacific islands: case study at Lakeba Island in Fiji. *Journal of Japan Society of Civil Engineers, Ser. B2 (Coastal Engineering)* **76**, I_1231–I_1236.
- UK Gov Marine Developments (2019) *Henderson Island Expedition*. Henderson Island Expedition – Marine developments (blog.gov.uk) (Accessed online 18 March 2021).
- UNEP (2017) *Coral Bleaching Futures – Downscaled Projections of Bleaching Conditions for the World's Coral Reefs, Implications of Climate Policy and Management Responses*. Nairobi: United Nations Environment Programme.
- UNEP-WCMC and IUCN (2021) Protected Planet. *Marine Protected Areas: The World Database on Protected Areas (WDPA)/The Global Database on Protected Areas Management Effectiveness (GD-PAME)*. Cambridge: UNEP-WCMC and IUCN. Available at www.protectedplanet.net (Accessed online 7 August 2021).
- Van Hooijdonk R, Maynard J, Tamelander J, Gove J, Ahmadi G, Raymundo L, Williams G, Heron SF and Planes S (2016) Local-scale projections of coral reef futures and implications of the Paris Agreement. *Nature* **6**, 1–8.
- Van Vuuren DP, Edmonds J, Kainuma M, Keywan R, Thomson A, Hibbard K, Hurtt GC, Kram T, Krey V, Lamarque J-F, Masui T, Meinshausen M, Nakicenovic N, Smith SJ and Rose SK (2011) The representative concentration pathways: an overview. *Climatic Change* **109**, 5.
- Vargas-Ángel B, Huntington B, Brainard RE, Venegas R, Oliver T, Barkley H and Cohen Brainard A (2019) El Niño-associated catastrophic coral mortality at Jarvis Island, central Equatorial Pacific. *Coral Reefs* **38**, 731–741.
- Venegas RM, Oliver T, Liu G, Heron SF, Clark SJ, Pomeroy N, Young C, Eakin CM and Brainard RE (2019) The rarity of depth refugia from coral bleaching heat stress in the Western and Central Pacific islands. *Scientific Reports* **9**, 19710.
- Vousdoukas MI, Mentaschi L, Voukouvalas E, Verlaan M, Jevrejeva S, Jackson LP and Feyen L (2018) Global probabilistic projections of extreme sea levels show intensification of coastal flood hazard. *Nature Communications* **9**, 1–12.
- Wagner T, Metzl N, Caffin M, Fin J, Helias Nunige S, Lefevre D, Lo Monaco C, Rougier G and Moutin T (2018) Carbonate system distribution, anthropogenic carbon and acidification in the Western Tropical South Pacific (OUTPACE 2015 transect). *Biogeosciences Discussions* **15**, 5221–5236. <https://doi.org/10.5194/bg-15-5221-2018>
- Wang XL and Swail VR (2006) Climate change signal and uncertainty in projections of ocean wave heights. *Climate Dynamics* **26**, 109–126.
- Weller E, Min SK, Cai W, Zwiers FW, Kim YH and Lee D (2016) Human-caused Indo-Pacific warm pool expansion. *Science Advances* **2**, e1501719.
- Wiedenmann J, D'Angelo C, Smith EG, Hunt AN, Legiret F-E, Postle AD and Achterberg EP (2013) Nutrient enrichment can increase the susceptibility of reef corals to bleaching. *Nature Climate Change* **3**, 160–164.
- Wilkinson CR (ed.) (2008) *Status of Coral Reefs of the World: 2008*. Townsville: Global Coral Reef Monitoring Network and Reef and Rainforest Research Centre, p. 296.
- Wood S, Paris CB, Ridgwell A and Hendy EJ (2014) Modelling dispersal and connectivity of broadcast spawning corals at the global scale. *Global Ecology and Biogeography* **23**, 1–11.
- Woodworth PL and Hibbert A (2015) Sea-level monitoring in the British overseas territories. *Journal of Operational Oceanography* **8**, 123–132.
- The World Bank (2016) *Climate and Disaster Resilience. Pacific Possible*. World Bank Group. Available at <http://pubdocs.worldbank.org/en/720371469614841726/PACIFIC-POSSIBLE-Climate.pdf>.
- Young IR, Zieger S and Babanin AV (2011) Global trends in wind speed and wave height. *Science (New York, N.Y.)* **332**, 451–455.
- Zhang X, Church J, Monselesan D and White N (2014) *Regional Sea Level Projection – Annual Time Series, 2006–2090 – 0.5 Degree Resolution*. V1. CSIRO. Data Collection. Available at <https://www.pacificclimatechange.net/document/regional-sea-level-projection-annual-time-series-2006-2090> (Accessed online 25 September 2020).