

## Women's iodine status and its determinants in an iodine-deficient area in the Kayes region, Mali

Liv E Torheim<sup>1,\*</sup>, Gry I Granli<sup>2</sup>, Cheick S Sidibé<sup>3</sup>, Abdel K Traoré<sup>4</sup> and Arne Oshaug<sup>1</sup>  
<sup>1</sup>Akershus University College, PO Box 423, N-2001 Lillestrøm, Norway; <sup>2</sup>Pediatric Ward, Rikshospitalet, Oslo, Norway; <sup>3</sup>Ecole Secondaire de la Santé, Bamako, Mali; <sup>4</sup>Faculté de Médecin, Pharmacie et d'Odonto-Stomatologie, Université du Mali, Bamako, Mali

Submitted 17 June 2004; Accepted 25 October 2004

### Abstract

**Objective:** To assess iodine status and its determinants in women of childbearing age in a rural area in the Kayes region, Mali, West Africa.

**Design:** Cross-sectional study where women's iodine status was indicated by urinary iodine concentration (UIC) and level of goitre based on palpation. Salt iodine was assessed semi-quantitatively at household level. Individual characteristics were collected using questionnaires.

**Setting:** Fifteen villages in a rural area in the Kayes region of Mali.

**Subjects:** Women aged 15–45 years ( $n = 423$ ).

**Results:** Median UIC was  $2.7 \mu\text{g dl}^{-1}$ , and only 6% of the women had adequate iodine status of  $\text{UIC} > 10 \mu\text{g dl}^{-1}$ . Most women (60%) had visible goitre, and only 9% were classified as without goitre. Only 39% of the households were using salt with any iodine, and level of knowledge about salt iodisation was low. Main determinants of UIC were breast-feeding and level of salt iodisation; currently breast-feeding women had lower UIC, and UIC increased with increasing level of iodine in household salt. Prevalence of goitre was lower in older women with higher body mass index.

**Conclusion:** The study indicates severe iodine deficiency in the study area. Urgent action is needed to improve the situation through enforcing salt iodisation legislation and increasing the level of knowledge about the importance of iodised salt in the population.

**Keywords**  
Iodine deficiency  
Iodised salt  
Urinary iodine concentration  
Women  
Human nutrition  
Mali  
Africa

Iodine deficiency still constitutes the single greatest cause of preventable brain damage and mental retardation<sup>1</sup>. Iodine deficiency disorders (IDD) *in utero* is recognised as a cause of poor mental and cognitive development<sup>2</sup>. All degrees of iodine deficiency affect the thyroid function of the mother and neonate, and the mental development of the child<sup>1</sup>. The resulting mental deficiency has an immediate effect on child learning capacity, women's health, the quality of life of communities and economic production<sup>3</sup>. Other consequences of iodine deficiency and subsequent inadequate thyroid hormone production include goitre, hypothyroidism, cretinism, congenital anomalies, increased perinatal mortality, stillbirths and abortions<sup>4</sup>. Much has been achieved in improving the coverage of households with iodised salt, which is the main intervention to address IDD. In developing countries approximately 66% of households are now using iodised salt<sup>5</sup>. Recent estimates by the World Health Organization (WHO) indicate that 54 countries are still affected by iodine deficiency and nearly 2 billion individuals worldwide are iodine-deficient<sup>6</sup>.

In Mali, salt iodisation was adopted as the preventive method for iodine deficiency in 1995, and the same year legislation on salt iodisation was passed<sup>7</sup>. Most of the salt is imported from Senegal, and all imported salt should be fortified with at least 25 g iodine per kg salt<sup>8</sup>. No nationwide survey of the IDD situation in Mali has been conducted. A Demographic Health Survey (DHS) in 2001 found that 74% of households were using iodised salt<sup>9</sup>. An assessment of the food and nutrition situation in Bafoulabé in the Kayes region was conducted in 1997 as part of a collaboration between researchers and a local non-government organisation. The survey registered cases of visible goitre only, and found a high prevalence in the area<sup>10</sup>. The present study was conducted to investigate further the iodine situation in the area, both the extent of iodine deficiency and determinants for planning suitable actions. Women were targeted since they constitute a group that is particularly vulnerable to IDD<sup>3</sup>.

The main objective of the present paper is to assess iodine status and its determinants in women of childbearing age in this rural area in the Kayes region of Mali.

\*Corresponding author. Email livelin.torheim@hiak.no

## Subjects and methods

### Study area

The study was done in Oussoubidiana development sector, Bafoulabé district, in the Kayes region of Western Mali. Oussoubidiana is a remote area of approximately 7500 inhabitants, where subsistence agriculture and raising livestock are the main livelihood activities<sup>11</sup>. The main ethnic group in this community is Kassonké. The most important foods are sorghum, maize, groundnuts and green leaves, and food insecurity is a seasonal concern<sup>11</sup>.

### Sample population

The target population was women of reproductive age (15–45 years) recruited from all 15 villages that constituted Oussoubidiana development sector. A sample size of 300 women was needed, derived from an assumed prevalence of iodine deficiency of 50% and a 95% confidence interval<sup>12</sup>. This corresponded to an estimated 160 households (including an estimated drop-out of 20%). The number of households to be selected in each of the 15 villages was determined based on the size of village. The households were randomly selected in each village, and all women aged 15–45 years in the selected households were included. The final number of study participants was 423 women recruited from 150 different households. Women's response rate was 92%, and the main reason for not participating was absence at the day of visit.

Characteristics of the study participants are shown in Table 1. Median age was 25 years and age was skewed to the lower range of 15–45 years. Most women (77%) had given birth at least once, and median parity was 3.

**Table 1** Background characteristics of the study participants. Women aged 15–45 years ( $n = 423$ ; Mali, 1999)

Characteristic	Value
Age (years)	25 (18, 33)*
Childbirths (number)†	3 (1, 6)*
None	97 (23)‡
1–2	101 (24)‡
3–4	86 (20)‡
≥ 5	138 (33)‡
Stillbirths (number)	
0	309 (73)‡
1	82 (19)‡
> 1	32 (8)‡
Currently pregnant	56 (13)‡
Currently breast-feeding	156 (37)‡
Height (m)	1.63 (1.60, 1.67)*
Weight (kg)	54.9 (50.7, 60.5)*
Body mass index ( $\text{kg m}^{-2}$ )§	20.4 (19.1, 22.1)*
< 18.5	53 (15)‡
18.5 to ≤ 25	291 (80)‡
> 25	21 (6)‡

\* Median (25th, 75th percentile).

† Includes live births and stillbirths.

‡  $n$  (%).

§  $n = 365$ , since body mass index was calculated only for non-pregnant women, and height measures were missing for two of these.

Twenty-seven per cent of the women said they had experienced at least one stillbirth. Thirteen per cent of the women were currently pregnant, whereas 37% were breast-feeding. No woman said she was both pregnant and breast-feeding. Body mass index (BMI) was in the lower range (median of  $20.4 \text{ kg m}^{-2}$ ), and 15% of the study participants were underweight ( $\text{BMI} < 18.5 \text{ kg m}^{-2}$ ).

### Data collection

Five fieldworkers were recruited locally. All were fluent in both the local language (Kassonké) and French, and had at least 3 years' higher education in health or social work. Two fieldworkers had experience from previous surveys. The fieldworkers underwent a week of theoretical and practical training prior to the data collection, and they participated in a pilot test of the methods and the subsequent modifications. During the study, two supervisors (G.I.G. and C.S.S.) checked all data collected daily for errors and completeness.

A pre-coded questionnaire was administered to the women by interview in the local language. It included questions on number of childbirths and number of stillbirths, and whether she was currently pregnant and/or breast-feeding. The women were weighed lightly clothed, using digital scales (Soehnle 7505, 100 g precision). Standing height was measured using a locally made stadiometer with 0.1 cm precision. The size of each woman's thyroid gland was visually inspected and palpated, and was graded according to the 1960 WHO criteria<sup>13</sup> with five categories of goitre severity. A sample of morning urine was obtained from each participant. The urine samples were kept frozen until analysed. Urinary iodine concentration (UIC) was measured using the Sandell–Kolthoff reaction<sup>14,15</sup>.

The household head, the traditional provider of salt, was asked whether the household salt was iodised, as well as its provenance. The household salt was tested for iodine content using a semi-quantitative test kit (Machine Build Industries, India). These are field test kits for testing salt fortified with potassium iodate ( $\text{KIO}_3$ ). Potassium iodate is the compound used for fortification of the salt imported from Senegal to Mali. The iodine concentration in the salt was determined visually from a colour chart with five codes: 0, 25, 50, 75 and above 100 parts of iodine per million (ppm).

Study permission was obtained from the Malian National Centre for Scientific and Technological Research, district authorities and village leaders. Each woman gave informed, oral consent to participate. The survey was conducted in November–December 1999.

### Data analysis

The five groups of goitre of the 1960 classification were collapsed into three groups that corresponded to the simplified classification suggested by WHO, the United Nations Children's Fund (UNICEF) and the International

Council for Control of Iodine Deficiency Disorders (ICCIDD) in 1994<sup>3,12</sup>: grade 0 remained the same (not palpable), grade 1A and 1B equalled the new grade 1 (palpable but not visible), whereas grade 2 to 3 became the new grade 2 (visible goitre when the neck is in normal position). The re-classification was done to ensure large enough groups in the data analysis.

The IDD indicators assessed (UIC, goitre and iodine content of household salt) were compared with WHO/ICCIDD/UNICEF epidemiological criteria for assessing the severity of IDD in a population<sup>3</sup>. According to these guidelines, median UIC below 2.0  $\mu\text{g dl}^{-1}$  indicates severe iodine deficiency, between 2.0 and 4.9  $\mu\text{g dl}^{-1}$  indicates moderate iodine deficiency, and between 5.0 and 9.9  $\mu\text{g dl}^{-1}$  shows mild iodine deficiency is present. Furthermore, a geographical area with a goitre rate above 30% is classified as severely iodine-deficient, between 20 and 29.9% as moderately iodine-deficient and between 5 and 19.9% as mildly iodine-deficient. One goal for sustainable elimination of IDD has been expressed as more than 90% of the households use adequately iodised salt, which has been defined as at least 15 ppm in populations which consume on average 10 g salt per day<sup>3</sup>.

Continuous variables are expressed as median and percentiles, and were compared by Kruskal–Wallis and Mann–Whitney tests or Spearman's rank correlation. Categorical variables were compared using the chi-square test. Linear regression was used to identify the determinants of UIC, using the square root of UIC to obtain a normal distribution. *P*-values of less than 0.05 were considered significant.

## Results

Median UIC of the study participants was 2.7  $\mu\text{g dl}^{-1}$  (Table 2). Most women were severely or moderately iodine-deficient (35% and 40%, respectively). Adequate iodine status of > 10  $\mu\text{g dl}^{-1}$  iodine/dl urine was found in only 6% of the sample. Table 2 also shows that only 9% of the women were classified as without goitre based on palpation, and most women (60%) had visible goitre (grade 2).

The majority of the households (61%) used salt with no iodine as tested with the semi-quantitative test kit (Table 3). Only 2% of the respondents said their household salt was iodised, and most households did not know whether their salt was iodised (69%). There was no relationship between the household's perception of whether the salt was iodised and the tested level of iodine in the household salt (data not shown).

Table 4 shows the association between the various characteristics of the study participants and their UIC. There was a positive association between the women's UIC and iodine level in household salt ( $P < 0.001$ ). Among those using salt without any detected iodine,

**Table 2** Iodine status of women ( $n = 423$ ; Mali, 1999)

Iodine status indicator	Value
Urinary iodine ( $\mu\text{g dl}^{-1}$ )*	2.7 (1.3, 4.9)†
< 2.0	140 (35)‡
2.0–4.9	170 (40)‡
5.0–9.9	78 (18)‡
$\geq 10.0$	26 (6)‡
Goitre – 1991 classification§	
Grade 0	37 (9)‡
Grade 1	131 (31)‡
Grade 2	255 (60)‡
Goitre – 1960 classification¶	
Grade 0	37 (9)‡
Grade 1A	30 (7)‡
Grade 1B	101 (24)‡
Grade 2	222 (52)‡
Grade 3	33 (8)‡

\* To convert to  $\mu\text{mol dl}^{-1}$ , multiply by 0.0079.

† Median (25th, 75th percentile).

‡ *n* (%).

§ Grade 0 – no palpable goitre; grade 1 – not visible but palpable goitre; grade 2 – visible goitre.

¶ Grade 0 – no palpable goitre; grade 1A – not visible but palpable goitre with neck in normal position; grade 1B – not visible but palpable goitre with neck in normal position, and visible with neck in extended position; grade 2 – visible and palpable goitre with neck in normal position, and goitre is not visible at a distance of more than 10 m; grade 3 – visible goitre at a distance of 10 m.

median UIC was 1.9  $\mu\text{g dl}^{-1}$ , below the WHO/ICCIDD/UNICEF cut-off for severe iodine deficiency<sup>3</sup>. The women from households using salt with 25 and 50 ppm iodine fell in the category of moderately iodine-deficient, with median UIC of 3.4 and 4.0  $\mu\text{g dl}^{-1}$ , respectively. Even those using salt with at least 75 ppm were mildly iodine-deficient, with median UIC of 7.0  $\mu\text{g dl}^{-1}$ . Among women from households using salt with any detected iodine (at least 25 ppm) only 13% had an adequate UIC above 10  $\mu\text{g dl}^{-1}$  (data not shown).

The women who were currently breast-feeding had significantly lower UIC compared with both those neither pregnant nor breast-feeding and those who were pregnant (median values of 2.3, 3.0 and 2.8  $\mu\text{g dl}^{-1}$ , respectively,  $P = 0.008$ ). There was no difference in UIC between pregnant and non-pregnant women. None of the other characteristics tested (parity, stillbirths, age and BMI) were found to have any association with UIC (Table 4).

To separate the effect of breast-feeding from the use of iodised salt, ordinary least-squares regression was used

**Table 3** Household use of iodised salt ( $n = 150$ ; Mali, 1999)

Variable	<i>n</i> (%)
Salt iodisation level (ppm)	
0	92 (61)
25	28 (19)
50	19 (13)
75	10 (7)
100	1 (0)
Answer to question on whether household salt was iodised	
Not iodised	44 (29)
Iodised	3 (2)
Did not know	103 (69)

**Table 4** Association between urinary iodine ( $\mu\text{g dl}^{-1}$ ) and background and nutritional characteristics ( $n = 423$ ; Mali, 1999)

	Median (25th, 75th percentile)
Iodine level in salt (ppm)	
0 ( $n = 249$ )	1.9 (0.9, 3.2)
25 ( $n = 70$ )	3.4 (2.7, 5.6)
50 ( $n = 73$ )	4.0 (2.6, 7.8)
$\geq 75^*$ ( $n = 31$ )	7.0 (3.7, 11.1)
<i>P</i> -value†	<0.001
Parity (number)	
0 ( $n = 97$ )	3.0 (1.9, 5.3)
1–2 ( $n = 101$ )	2.6 (1.3, 5.1)
3–4 ( $n = 86$ )	2.7 (0.9, 4.9)
$\geq 5$ ( $n = 138$ )	2.6 (1.3, 4.5)
<i>P</i> -value†	0.2
Stillbirths (number)	
0 ( $n = 309$ )	2.7 (1.4, 5.0)
1 ( $n = 82$ )	2.9 (1.5, 5.2)
$\geq 2$ ( $n = 32$ )	2.2 (1.1, 3.6)
<i>P</i> -value†	0.1
Current reproductive status	
Neither pregnant nor breast-feeding ( $n = 212$ )	3.0 (1.6, 5.2)
Pregnant ( $n = 56$ )	2.8 (1.6, 5.6)
Breast-feeding ( $n = 156$ )	2.3 (1.0, 3.8)
<i>P</i> -value†	0.008
Age (years)	
<i>r</i>	–0.04
<i>P</i> -value	0.4
Body mass index ( $\text{kg m}^{-2}$ )‡	
<i>r</i>	0.00
<i>P</i> -value	1.0

*r* – Spearman's correlation coefficient.

\* Women from households using salt with 75 ppm ( $n = 29$ ) and with 100 ppm ( $n = 2$ ) added.

† Differences tested with Kruskal–Wallis one-way analysis of variance.

‡ Includes only non-pregnant women.

(Table 5). Both variables had significant association with UIC when entered into the same model, which explained 24% of the variation. Figure 1 illustrates the impact of iodisation level of household salt and breast-feeding on UIC.

The same characteristics, as well as UIC, were analysed in relation to goitre (Table 6). There was a tendency towards a lower prevalence of visible goitre in the groups with higher UIC ( $P = 0.07$ ). Age and BMI were found to be associated with goitre; goitre prevalence was highest among the youngest and thinnest groups of women. Figure 2 illustrates that the association between BMI and goitre existed only in the highest age group. Most of those

without goitre were in the highest age group and had the highest BMI. The association between age and goitre was similar, and it existed only in the highest BMI tertile (data not shown).

Parity was also found to be associated with goitre. Nulliparous women had the highest prevalence of goitre, with 98% classified as goitrous grade 1 or 2, whereas among those with at least five children, 84% were classified as having goitre ( $P = 0.003$ ). However, this association did not remain significant when controlling for age. No relationships were found between goitre and iodine level in salt, current reproductive status or number of stillbirths.

## Discussion

Our study showed widespread iodine deficiency among women in this area of rural Western Mali. Median UIC was  $2.7 \mu\text{g dl}^{-1}$  and only 6% had adequate UIC, i.e. above  $10 \mu\text{g dl}^{-1}$ . As many as 75% of the women had UIC below  $5 \mu\text{g dl}^{-1}$ , being moderately to severely iodine-deficient according to the epidemiological criteria set by WHO/IC-CIDD/UNICEF<sup>5</sup>. The overall prevalence of goitre of 91% was extremely high. It greatly exceeded the limit of 10% where IDD is considered a public health problem<sup>5</sup>. Only 39% of the households used iodised salt (at least 25 ppm), and knowledge about iodised salt was low.

All three IDD indicators investigated show that Oussoubidiana is a pocket of extreme iodine deficiency where specific measures are needed. Recent studies from other parts of Mali give a less grim picture of the IDD situation in the country. A study was done in 1999 by ThyroMobil (a van equipped with ultrasound for measurement of thyroid volume, a computer and a deep freezer) and included samples of schoolchildren in nine sites around Segou in Mali<sup>16</sup>. The study found a median UIC of  $20.3 \mu\text{g dl}^{-1}$  and 66% of the sample had UIC above  $10 \mu\text{g dl}^{-1}$ . Total goitre rate was 13%, as assessed by ultrasonography. A study of adolescent girls (10–15 years of age) in the region of Segou<sup>17</sup> found a mean UIC of  $9.3 \mu\text{g dl}^{-1}$  and 33% had UIC above  $10 \mu\text{g dl}^{-1}$ . In the DHS of 2001<sup>9</sup>, overall 74% of households in Mali were found to be using iodised salt, and in the region of Kayes the prevalence was 69%. There is no evidence pointing

**Table 5** Determinants of urinary iodine\* ( $n = 423$ ; Mali, 1999)


Variable	Unadjusted effect (95% confidence interval)	<i>P</i> -value	Adjusted effect† (95% confidence interval)	<i>P</i> -value
Constant			1.25 (1.10, 1.40)	<0.001
Iodine level in salt‡	0.45 (0.38, 0.55)	<0.001	0.47 (0.39, 0.55)	<0.001
Currently breast-feeding§	0.21 (0.02, 0.40)	0.032	0.23 (0.07, 0.40)	0.006
$R^2$			0.24	

\* Square root of urinary iodine concentration ( $\mu\text{g dl}^{-1}$ ).


† Adjusted for the other variable in the table in multiple regression analysis.

‡ Categories of iodine level in salt: 0 = 0 ppm; 1 = 25 ppm; 2 = 50 ppm; 3 =  $\geq 75$  ppm.

§ Breastfeeding dummy: 0 = yes; 1 = no.



**Fig. 1** Comparison of (square-root) urinary iodine concentration (UIC) in breast-feeding and non-breast-feeding women according to level of household salt iodisation as measured by rapid test kit ( $n = 423$ ; Mali, 1999)



**Fig. 2** Comparison of (ln-transformed) body mass index (BMI) between women with no goitre (grade 0), palpable goitre (grade 1) and visible goitre (grade 2) in three age groups ( $n = 423$ ; Mali, 1999)

towards the Kayes region as being a particularly severely affected region in Mali. Much of the focus on iodine deficiency has been in the central regions, like Segou and Mopti.

Oussoubidiana is a remote and relatively inaccessible area of the Kayes region, which might contribute to the severe IDD situation. It is unclear whether the salt

reaching the area was not iodised at all, or whether there had been a loss of iodine due to storage over longer periods. High losses of iodine have been shown from salt packed in high-density polyethylene bags<sup>18</sup>. This was the packaging material mainly used at storage and retail level in the study area. In their study from the plains of Nepal, Schulze *et al.*<sup>19</sup> showed that urinary iodine of pregnant

**Table 6** Association between goitre prevalence (%) and background and nutritional characteristics ( $n = 423$ ; Mali, 1999)

	Grade 0 ( $n = 31$ )	Grade 1 ( $n = 131$ )	Grade 2 ( $n = 255$ )	<i>P</i> -value*
Iodine level in salt (ppm)				
0 ( $n = 249$ )	10	29	61	0.5
> 0 ( $n = 174$ )	7	34	59	
Urinary iodine concentration ( $\mu\text{g dl}^{-1}$ )				
< 2 ( $n = 149$ )	7	31	62	0.07
2–4.9 ( $n = 170$ )	10	25	65	
5–9.9 ( $n = 78$ )	8	37	55	
$\geq 10$ ( $n = 26$ )	8	54	38	
Parity (number)				
0 ( $n = 97$ )	2	36	62	0.003
1–2 ( $n = 101$ )	9	34	57	
3–5 ( $n = 86$ )	5	36	59	
$\geq 5$ ( $n = 138$ )	16	22	62	
Stillbirths (number)				
0 ( $n = 309$ )	7	32	61	0.3
1 ( $n = 82$ )	12	29	59	
$\geq 2$ ( $n = 32$ )	16	22	62	
Current reproductive status				
Not pregnant or breast-feeding ( $n = 212$ )	9	30	61	0.9
Pregnant ( $n = 55$ )	7	29	64	
Breast-feeding ( $n = 156$ )	8	33	58	
Age (years, tertiles)				
15–20 ( $n = 142$ )	2	34	63	<0.001
21–29 ( $n = 131$ )	6	36	58	
30–45 ( $n = 150$ )	17	23	59	
Body mass index ( $\text{kg m}^{-2}$ , tertiles)†				
14.0–19.6 ( $n = 121$ )	7	26	67	<0.001
19.7–21.4 ( $n = 122$ )	2	43	55	
21.5–35.4 ( $n = 122$ )	19	25	57	

Grade 0 – no palpable goitre; grade 1 – not visible but palpable goitre; grade 2 – visible goitre.

\* Differences tested with chi-squared test.

† Includes only non-pregnant women.

and lactating women and salt iodine covaried markedly by season, and were highest during hot and dry pre-monsoon months and lowest during and following the humid monsoon season. Our study was conducted in a cooler period some months after the rainy season, with a climate that should not have had adverse effects on the iodine content of the salt. Iodine may nevertheless have been lost from salt during transportation or storage.

Spot checks by the study team of salt at storage and retail level found both non-iodised salt and salt with low iodine (ppm) content in bags marked as being iodised. There were also bags not marked as being iodised, containing non-iodised salt. Current Malian legislation prohibits import and distribution of non-iodised salt. Although the study was done in 1999, no concrete action has been taken at national level to improve the situation. These results therefore point to a need for urgent control and monitoring of iodine content in salt at production, import and resale points. This should be based on enforcement of the existing legislation. An in-depth study of the origin of salt used in this area, mapping of salt importers and training of these are necessary to plan and implement an effective monitoring system. Efforts to ensure adequate iodine levels in salt at all points should be combined with campaigns to increase population awareness about the importance of iodine and create public demand for iodised salt.

#### **Determinants of UIC**

The main determinant of UIC was the level of iodine in the household salt. However, despite the positive correlation between UIC and content of iodine in household salt, most of the women using salt with iodine level of 25 ppm or above still had UIC below  $10 \mu\text{g dl}^{-1}$  (77%). These results indicate that the recommended iodine content of 15 ppm at household level may not be sufficient for sustaining an adequate iodine status in women of fertile age in this area. Dietary surveys conducted in the area have indicated a daily salt intake of around  $10 \text{ mg}^{20,21}$ , which is the basis for the current level of iodine fortification. Possible explanations for the lower than expected UIC in women from households using iodised salt could be consumption of food made with salt other than that tested in the household or loss of iodine during cooking. Studies have shown losses of up to 70% of salt iodine during cooking<sup>22</sup>. Imprecise or inaccurate measurement of iodine level in the household salt must also be considered. An evaluation of rapid test kits showed that they might lead to over-estimation of the iodine content in salt<sup>23</sup>. Nevertheless, our results demonstrate the necessity of assessing UIC, and not only salt iodine content, in monitoring IDD.

Pregnancy was not found to be associated with UIC, unlike what was found in a study from another iodine-deficient area in Sudan<sup>24</sup>. Increased thyroid hormone requirements<sup>25</sup> and increased iodine needs for the developing foetus are combined with an increased renal

clearing of iodine during pregnancy<sup>26</sup>. It is therefore not evident that pregnancy will result in reduced UIC. Brander *et al.*<sup>27</sup> reported, in a study from Switzerland, significantly higher UIC in pregnant women in their first trimester compared with controls who were mildly iodine-deficient. Schulze *et al.*<sup>19</sup> found higher UIC in women during pregnancy compared with the same subjects during lactation in a moderately iodine-deficient area in Nepal.

Breast-feeding women had lower UIC than non-breast-feeding women, but only among those using salt with no iodine or low levels of iodine (0–25 ppm). Reduced iodine status during breast-feeding has been found also in other studies<sup>19,24</sup>, and is probably partly a reflection of the iodine excretion in breast milk. The iodine concentration of human milk varies widely linked to maternal iodine intake<sup>28</sup>. Undoubtedly, the low iodine status of the pregnant and breast-feeding women in this study may lead to a negative impact on the mental development of the foetus and the neonate<sup>1</sup>.

#### **Determinants of goitre**

Goitre prevalence decreased with age and parity. Regarding the toll repeated pregnancies and breast-feeding periods take on women's iodine status, this was surprising. These results were contradictory to other studies that have shown an increasing thyroid enlargement with age in women in areas of severe<sup>29</sup> or moderate to mild iodine deficiency<sup>30,31</sup>. Parity has also been related to increased thyroid volume in moderately iodine-deficient areas<sup>30,32</sup>. It has been suggested that there is no effect of age *per se*, but that the relationship between thyroid size and age is solely due to parity<sup>30</sup>. The higher levels of goitre found in the nulliparous, as compared with the multiparous, in our study might reflect the young age of the first group (81% were less than 20 years old). These young women might have had an increased need for iodine due to increased production of thyroid hormones during puberty.

Goitre status was not related to either current pregnancy or breast-feeding in our study. This was unexpected, since pregnancy has been reported to bring about goitre even in moderately iodine-deficient areas of Europe<sup>33,34</sup>. Goitres formed during gestation may regress only partially after parturition<sup>35</sup>. It has therefore been suggested that pregnancy represents an environmental factor that may help explain the higher prevalence of goitre and thyroid disorders in women compared with men<sup>36</sup>.

There was a negative association between goitre prevalence and BMI. Some studies have indicated that malnutrition might in itself lead to abnormal thyroid function<sup>37</sup>. In our study, the negative relationship between BMI and goitre was found only among the oldest women. One explanation for this observation could be that a high BMI in older women reflected that they had not given birth for some time. This could have allowed iodine and energy stores depleted owing to the strains of pregnancy and

breast-feeding to be restored. This study did not give any information about when the women had last given birth; however, fertility decreased with age.

Misclassification is a major problem of assessment of thyroid enlargement by palpation<sup>38</sup> and cannot be ruled out as an explanation for our results. A systematic misclassification would give a type 1 error (establish a non-existing association), whereas random misclassification would give a type 2 error (failing to observe an existing association)<sup>39</sup>. It is possible that younger and thinner women have wrongly been classified as goitrous to a larger extent than older women with higher BMI.

### Conclusion

In conclusion, this study has shown a situation of severe iodine deficiency among women of childbearing age in this area of rural Kayes, Western Mali. Inadequate salt iodisation combined with increased need for iodine during pregnancy and lactation were the main determinants. Urgent action is needed to improve the situation through increased access to and knowledge about iodised salt.

### Acknowledgements

Our thanks go to the women who patiently answered our questions and inquiries. Thanks also to the fieldworkers who did a painstaking job with great efforts. We greatly appreciate the support of the Stromme Foundation in Bamako and of AIDEB in Bafoulabé. This study was funded by the Stromme Foundation. We also thank UNICEF Mali for providing the semi-quantitative test kits to test for iodine in salt.

### References

- Delange F. The role of iodine in brain development. *Proceedings of the Nutrition Society* 2000; **59**: 75–9.
- Pharoah POD, Connolly KJ. Iodine deficiency in Papua New Guinea. In: Stanbury JB, ed. *The Damaged Brain of Iodine Deficiency*. New York: Cognizant Communication Corporation, 1994; 299–305.
- World Health Organization (WHO)/International Council for Control of Iodine Deficiency Disorders/United Nations Children's Fund. *Assessment of Iodine Deficiency Disorders and Monitoring Their Elimination – A Guide for Programme Managers*. Geneva: WHO, 2001.
- Hetzel BS, Dunn JT, Stanbury JB. *The Prevention and Control of Iodine Deficiency Disorders*. Amsterdam: Elsevier, 1987.
- United Nations Children's Fund (UNICEF). *The State of the World's Children 2004*. New York: UNICEF, 2003.
- United Nations Standing Committee on Nutrition (SCN). *5th Report on the World Nutrition Situation*. Geneva: SCN, 2004.
- World Health Organization (WHO)/United Nations Children's Fund/International Council for Control of Iodine Deficiency Disorders. *Conference on Sustainable Elimination of Iodine Deficiency Disorders in Africa by the Year 2000*. Geneva: WHO, 1998.
- Global Micronutrient Project. *Iodine Situation in Mali* [online]. New Orleans, LA: University of Tulane. Available at <http://www.tulane.edu/~internut/Countries/Mali/malixx.html>. Accessed April 2004.
- Cellule de Planification et de Statistique du Ministère de la Santé (CPS/MS), Direction Nationale de la Statistique et de l'Informatique (DNSI) and ORC Macro. *Enquête Démographique et de Santé au Mali 2001*. Calverton, MD: CPS/MS, DNSI and ORC Macro, 2001.
- Torheim LE. *Assessing Dietary Intake in Rural Mali – Challenges when People eat from Shared Plates*. Oslo: Faculty of Medicine, University of Oslo, 2003.
- Torheim LE, Ouattara F, Diarra MM, Thiam FD, Barikmo I, Hatloy A, et al. Nutrient adequacy and dietary diversity in rural Mali: association and determinants. *European Journal of Clinical Nutrition* 2004; **58**: 594–604.
- World Health Organization (WHO)/United Nations Children's Fund/International Council for Control of Iodine Deficiency Disorders. *Indicators for Assessing Iodine Deficiency Disorders and Their Control through Salt Iodization*. Geneva: WHO, 1994.
- Perez C, Schrimshaw NS, Munoz JA. Technique of endemic goitre surveys. In: *Endemic Goiter*. World Health Organization (WHO) Monograph Series No. 44. Geneva: WHO, 1960; 369–83.
- Dunn JT, Crutchfield HE, Gutekunst R, Dunn AD. *Methods for Measuring Iodine in Urine*. The Netherlands: International Council for Control of Iodine Deficiency Disorders, 1993.
- Dunn JT, Crutchfield HE, Gutekunst R, Dunn AD. Two simple methods for measuring iodine in urine. *Thyroid* 1993; **3**: 119–23.
- Kibambe N, Ouedraogo A, Acakpo A, Salami M. *Projet Thyromobil en Afrique de L'Ouest (première phase). Rapport Final*. Lome: Conseil International de Lutte contre les Troubles Dus à la Carence en Iode/Organisation Mondiale de la Santé/Centre Régional de Recherche en Alimentation et Nutrition/Fonds des Nations Unies pour l'Enfance, 2000.
- Pawloski LR, Shier NW, Fernandez XE, Jamison PL. Examination of urinary iodine status from a sample of Malian adolescent girls. *African Journal of Food, Agriculture, Nutrition and Development* 2003; **3**: 20–8.
- Diosady LL, Alberti JO, Venkatesh Mannar MG, FitzGerald S. Stability of iodine in iodised salt used for correction of iodine-deficiency disorders II. *Food and Nutrition Bulletin* 1998; **19**: 240–9.
- Schulze KJ, West KP Jr, Gautschi LA, Dreyfuss ML, LeClerq SC, Dahal BR, et al. Seasonality in urinary and household salt iodine content among pregnant and lactating women of the plains of Nepal. *European Journal of Clinical Nutrition* 2003; **57**: 969–76.
- Torheim LE, Barikmo I, Hatloy A, Diakité M, Solvoll K, Diarra MM, et al. Validation of a quantitative food-frequency questionnaire for use in Western Mali. *Public Health Nutrition* 2001; **4**: 1267–77.
- Parr CL, Barikmo I, Torheim LE, Ouattara F, Kaloga A, Oshaug A. Validation of the second version of a quantitative food-frequency questionnaire for use in Western Mali. *Public Health Nutrition* 2002; **5**: 769–81.
- Verma M, Raghuvanshi RS. Dietary iodine intake and prevalence of iodine deficiency. *Journal of Nutritional and Environmental Medicine* 2001; **11**: 175–80.
- Pandav CS, Arora NK, Krishnan A, Sankar R, Pandav S, Karmarkar MG. Validation of spot-testing kits to determine iodine content in salt. *Bulletin of the World Health Organization* 2000; **78**: 975–80.
- Eltom A, Eltom M, Elnagar B, Elgaber M, Gebre-Medhin M. Changes in iodine metabolism during late pregnancy and lactation: a longitudinal study among Sudanese women. *European Journal of Clinical Nutrition* 2000; **54**: 429–33.
- Glinoe D. Feto-maternal repercussions of iodine deficiency during pregnancy. An update. *Annales d'Endocrinologie* 2003; **64**: 37–44.

- 26 Dunn JT, Delange F. Damaged reproduction: the most important consequence of iodine deficiency. *Journal of Clinical Endocrinology and Metabolism* 2001; **86**: 2360–3.
- 27 Brander L, Als C, Buess H, Haldimann F, Harder M, Hanggi W, *et al.* Urinary iodine concentration during pregnancy in an area of unstable dietary iodine intake in Switzerland. *Journal of Endocrinological Investigation* 2003; **26**: 389–96.
- 28 Dorea JG. Iodine nutrition and breast feeding. *Journal of Trace Elements in Medicine and Biology* 2002; **16**: 202–20.
- 29 Adesunkanmi AR, Makinde ON. Goitre prevalence in pregnant women attending antenatal clinic in a teaching hospital. *Journal of Obstetrics and Gynaecology* 2003; **32**: 156–9.
- 30 Rotondi M, Amato G, Biondi B, Mazziotti G, Del Buono A, Rotonda Nicchio M, *et al.* Parity as a thyroid size-determining factor in areas with moderate iodine deficiency. *Journal of Clinical Endocrinology and Metabolism* 2000; **85**: 4534–7.
- 31 Knudsen N, Bülow I, Jørgensen T, Laurberg P, Ovesen L, Perrild H. Goitre prevalence and thyroid abnormalities at ultrasonography: a comparative epidemiological study in two regions with slightly different iodine status. *Clinical Endocrinology* 2000; **53**: 479–85.
- 32 Knudsen N, Bülow I, Laurberg P, Ovesen L, Perrild H, Jørgensen T. Parity is associated with increased thyroid volume solely among smokers in an area with moderate to mild iodine deficiency. *European Journal of Endocrinology* 2002; **146**: 39–43.
- 33 Smyth PP, Hetherington AM, Smith DF, Radcliff M, O’Herlihy C. Maternal iodine status and thyroid volume during pregnancy: correlation with neonatal iodine intake. *Journal of Clinical Endocrinology and Metabolism* 1997; **82**: 2840–3.
- 34 Rasmussen NG, Hornnes PJ, Hegedus L. Ultrasonographically determined thyroid size in pregnancy and post partum: the goitrogenic effect of pregnancy. *American Journal of Obstetrics and Gynecology* 1989; **160**: 1216–20.
- 35 Glinoe D, Lemone M, Bourdoux P, De Nayer P, Delange F, Kinthaert J, *et al.* Partial reversibility during late postpartum of thyroid abnormalities associated with pregnancy. *Journal of Clinical Endocrinology and Metabolism* 1992; **74**: 453–7.
- 36 Glinoe D. Pregnancy and iodine. *Thyroid* 2001; **11**: 471–81.
- 37 Gaitan JE, Mayoral LG, Gaitan E. Defective thyroïdal iodine concentration in protein–calorie malnutrition. *Journal of Clinical Endocrinology and Metabolism* 1983; **57**: 327–33.
- 38 Tonglet R, Bourdoux P, Dramaix M, Hennart P, Ermans AM. Interobserver variation in the assessment of thyroid enlargement: a pitfall in surveys of the prevalence of endemic goitre. *Food and Nutrition Bulletin* 1994; **15**: 64–70.
- 39 Altman DG. *Practical Statistics for Medical Research*. London: Chapman & Hall, 1997.